VERSION 2

APR 02, 2019

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.zpuf5nw

External link:

<https://doi.org/10.1371/journal.pntd.0007659>

Protocol Citation: Bo Yi, Yi Chen, Xiao Ma, Haibin Wang, Rong Wang, Keqin Ding, Lei Xie, Dongliang Zhang, Shuli Jiao, Xuying Lao, Yi-Chen Chiang, Yanhua Su, Benhua Zhao, Guozhang Xu, Tianmu Chen 2019. E gene amplification. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.zpuf5nw>

MANUSCRIPT CITATION:

Yi B, Chen Y, Ma X, Rui J, Cui J, Wang H, Li J, Chan S, Wang R, Ding K, Xie L, Zhang D, Jiao S, Lao X, Chiang Y, Su Y, Zhao B, Xu G, Chen T (2019) Incidence dynamics and investigation of key interventions in a dengue outbreak in Ningbo City, China. PLoS Negl Trop Dis 13(8): e0007659. doi: [10.1371/journal.pntd.0007659](https://doi.org/10.1371/journal.pntd.0007659)

E gene amplification V.2

PLOS Neglected Tropical Diseases

Bo Yi¹, Yi Chen¹, Xiao Ma¹, Wang², Haibin Wang¹, Rong Wang¹, Keqin Ding¹, Lei Xie¹, Dongliang Zhang¹, Shuli Jiao¹, Xuying Lao¹, Yanhua Guozhang Yi-Chen Chiang³, Su³, Benhua Zhao³, Xu¹, Tianmu Chen³

¹Ningbo Municipal Center for Disease Control and Prevention ;

²Haishu District Center for Disease Control and Prevention ;

³State Key Laboratory of Molecular Vaccinology and Molecular Diagnostics, School of Public Health, Xiamen University

Tianmu Chen

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Apr 02, 2019

Last Modified: Apr 02, 2019

PROTOCOL integer ID:
21972

- 1 The dengue virus strains obtained from cell culture were extracted by TGuide S32 magnetic bead method DNA/RNA extraction kit of Tiangen Biochemical Technology (Beijing) Co., Ltd. and amplified by one step RNA PCR Kit (Code No: DRR057A) reagent of Bao Bioengineering (Dalian) Co., Ltd.
- 2 The full sequence of E gene amplified by primer sequence was D1 (E) F: CAA GAA CCG AAA CA / GT GGA TGT C; D1 (E) R: GGC TGA TCG AAT TCC ACA CAC, and the length of amplified fragment was 1849 bp.
- 3 Reaction conditions: reverse transcription at 50 °C 30 minutes, reverse transcription at 94°C 2 minutes, reverse transcription at 94°C 30 seconds, reverse transcription at 52°C 30 seconds, reverse transcription at 72°C 2 minutes, 40 cycles and extension at 72°C. The amplified product was 5 µL, and 1.5% agarose gel electrophoresis was used to confirm the reaction product according to Marker position.