FEB 26, 2021

Efficacy, safety and complications of autologous fat grafting to the eyelids and periorbital area: A systematic review and meta-analysis

Fan Yang¹

¹Department of Plastic Surgery and Burns, Tangdu Hospital, The Fourth Military Medical University, Shaanxi, China

crystalangfan

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bstqnmw

Protocol Citation: Fan Yang 2021. Efficacy, safety and complications of autologous fat grafting to the eyelids and periorbital area: A systematic review and meta-analysis. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bstqnmw>

MANUSCRIPT CITATION:
References 1. Coleman SR. Longterm survival of fat transplants controlled demonstrations. Aesthetic Plastic Surgery. 1995;19:421-5. 2. Konn M. Structural fat grafting. British Journal of Surgery. 2005;92(5):657-. doi: 10.1002/bjs.4962. 3. Ramil ME. Fat Grafting in the Hollow Upper Eyelids and Volumetric Upper Blepharoplasty. Plast Reconstr Surg. 2017;140(5):889-97. Epub 2017/07/29. doi: 10.1097/PRS.0000000000003758. PubMed PMID: 28753160. 4. Lee W, Kwon SB, Oh SK, Yang EJ. Correction of sunken upper eyelid with orbital fat transposition flap and dermofat graft. J Plast Reconstr Aesthet Surg. 2017;70(12):1768-75. Epub 2017/06/18. doi: 10.1016/j.bjps.2017.05.003. PubMed PMID: 28619482. 5. Lin TM, Lin TY, Huang YH, Hsieh TY, Chou CK, Takahashi

DISCLAIMER

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to protocols.io is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with protocols.io, can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

H, et al. Fat Grafting for Recontouring Sunken Upper Eyelids With Multiple Folds in Asians-Novel Mechanism for Neoformation of Double Eyelid Crease. *Annals of plastic surgery*. 2016;76(4):371-5. Epub 2015/12/19. doi: 10.1097/sap.0000000000000668. PubMed PMID: 26678103. 6. Lin TM, Lin TY, Chou CK, Lai CS, Lin SD. Application of microautologous fat transplantation in the correction of sunken upper eyelid. *Plast Reconstr Surg Glob Open*. 2014;2(11):e259. Epub 2014/12/17. doi: 10.1097/GOX.0000000000000141. PubMed PMID: 25506542; PubMed Central PMCID: PMC4255902. 7. Yoonho Lee MD, Ph.D., Sungtaek Kwon, M.D., Ph.D., and Kun Hwang, M.D., Ph.D. Correction of Sunken and/or Multiply Folded Upper Eyelid by Fascia-Fat Graft. *Plast Reconstr Surg*. 2001;107(1):15-9. 8. Tonnard PL, Verpaele AM, Zeltzer AA. Augmentation blepharoplasty: a review of 500 consecutive patients. *Aesthetic surgery journal*. 2013;33(3):341-52. Epub 2013/03/22. doi: 10.1177/1090820x13478966. PubMed PMID: 23515379. 9. Rehman J, Traktuev D, Li J, Merfeld-Clauss S, Temm-Grove CJ, Bovenkerk JE, et al. Secretion of angiogenic and antiapoptotic factors by human adipose stromal cells. *Circulation*. 2004;109(10):1292-8. Epub 2004/03/03. doi: 10.1161/01.CIR.0000121425.42966.F1. PubMed PMID: 14993122. 10. Varghese J, Griffin M, Mosahebi A, Butler P. Systematic review of patient factors affecting adipose stem cell viability and function: implications for regenerative therapy. *Stem Cell Res Ther*. 2017;8(1):45. Epub 2017/03/01. doi: 10.1186/s13287-017-0483-8. PubMed PMID: 28241882; PubMed Central PMCID: PMC5329955. 11. Trepsat F. Periorbital rejuvenation combining fat grafting and blepharoplasties. *Aesthetic Plast Surg*

ABSTRACT

Background: In recent years, autologous fat grafting (AFG), also known as fat transfer or lipofilling, has been widely performed for periorbital rejuvenation and defect correction, although the evidence regarding its efficacy and safety is still lacking. Besides, with respect to the periorbital region, it is invariably the earliest appearance area of the facial aging phenomenon. Therefore, a systematic review and meta-analysis is needed to evaluate the efficacy and safety of this technique. **Methods:** A literature search was performed in PubMed, Embase, and the Cochrane library databases on November 11, 2020, adhering to the PRISMA guidelines, to identify all relevant articles. Then, a data extraction and standardization process was performed to assess all outcome data. Ultimately, the data were assessed using a random effects regression model with comprehensive meta-analysis software. **Results:** Thirty-six studies consisting of four cohorts and 32 case series with a total of 3997 cases were included. The mean follow-up was 1.5 years. Meta-analysis revealed a relatively high satisfaction rate of 93.7% (95% CI, 89.6%–96.2%). Frequent complications in 3997 patients receiving AFG were edema, chemosis, and contour irregularity, with an overall complication rate of 7.7% (95% CI, 4.3%–13.3%). **Conclusion:** This systematic review and meta-analysis showed that AFG for rejuvenation of eyelids and periorbital area provided a high satisfaction rate and did not result in severe complications. Therefore, AFG might be performed safely for periorbital rejuvenation and reconstruction.

GUIDELINES

The protocol was written according to the Preferred Reporting Items for Systematic Review and Meta-Analysis (PRISMA)

Aesthetic Plast Surg.
2003;27(4):243-53. Epub
2004/04/03. doi:
10.1007/s00266-003-2126-
y. PubMed PMID: 15058544.
12. Boureaux E, Chaput B,
Bannani S, Herlin C, De Runz
A, Carloni R, et al. Eyelid fat
grafting: Indications, operative
technique and complications;
a systematic review. *Journal of
cranio-maxillo-facial surgery* :
official publication of the
European Association for
Cranio-Maxillo-Facial Surgery.
2016;44(4):374-80. Epub
2016/02/18. doi:
10.1016/j.jcms.2015.12.013.
PubMed PMID: 26880013. 13.
Maamari RN, Massry GG,
Holds JB. Complications
Associated with Fat Grafting
to the Lower Eyelid. *Facial
plastic surgery clinics of North
America*. 2019;27(4):435-41.
Epub 2019/10/08. doi:
10.1016/j.fsc.2019.07.001.
PubMed PMID: 31587763. 14.
Shue S, Kurlander DE,
Guyuron B. Fat Injection: A
Systematic Review of Injection
Volumes by Facial Subunit.
Aesthetic plastic surgery.
2018;42(5):1261-70. Epub
2017/08/10. doi:
10.1007/s00266-017-0936-
6. PubMed PMID: 28791455.
15. Cetinkaya A, Devoto MH.
Periocular fat grafting:
indications and techniques.
Curr Opin Ophthalmol.
2013;24(5):494-9. Epub
2013/08/09. doi:
10.1097/ICU.0b013e328363
4841. PubMed PMID:
23925063. 16. Knobloch K,
Yoon U, Vogt PM. Preferred
reporting items for systematic
reviews and meta-analyses
(PRISMA) statement and
publication bias. *J
Craniofac Surg*.
2011;39(2):91-2. Epub
2010/12/15. doi:
10.1016/j.jcms.2010.11.001.
PubMed PMID: 21145753. 17.
Pelle-Ceravolo M, Angelini M.
Properly Diluted Fat (PDF): An
Easy and Safe Approach to
Periocular Fat Grafting.
Aesthetic surgery journal.
2019. Epub 2019/02/13. doi:
10.1093/asj/sjz039. PubMed
PMID: 30753276. 18.
Essuman VA, Tagoe NN,
Ndanu TA, Ntim-Amponsah
CT. Dermis-fat grafts and

MATERIALS

Search strategy

The research objectives were to identify, assess, and synthesize the evidence examining the efficacy and safety of AFG in the periocular area. This review was performed in accordance with the PRISMA guidelines. This comprehensive, reproducible, and electronic search was performed via the combination of PubMed, Embase, and Cochrane Library databases. The following keywords were used: [("fat grafting" OR "lipograft" OR "lipoinjection" OR "lipotransfer" OR "fat transfer" OR "fat transplant" OR "lipostructure" OR "lipofilling" OR "fat injection" OR "lipomodeling" OR "fat transplantation") AND ("eyelid" OR "periocular")]. A systematic database search was carried out before November 11, 2020. There were no restrictions with respect to language.

Inclusion and exclusion criteria

Selected studies met the following criteria: (1) clinical trials of all designs, from the highest level of evidence from randomized trials (if available) to prospective or retrospective observational studies (case series: at least five cases) involving patients receiving AFG for periorbital rejuvenation and reconstruction; (2) the treatment used for periorbital rejuvenation and reconstruction was stated clearly; (3) the study stated the concrete data of postoperative effects; (4) studies with complete follow-up (at least 3 months). Exclusion criteria were as follows: (1) patients with a history of other eyelid surgery or treatment; (2) reviews, letters, commentaries, reply, discussion, and so on; (3) studies with incomplete or ambiguous or overlapped data; (4) studies not related to the objective of this review.

Data collection

Two independent reviewers scrutinized the titles, abstracts, and full text of the retrieved articles. If there was any disagreement between the two reviewers, another independent investigator was consulted to reach a consensus. Moreover, a blinded method was used to ensure quality. Data extracted from the eligible articles included the following parts: authors, date of publication, place of study, number of patients, ages of patients, indications, AFG techniques, follow-up time, study design, evidence level, complications, anesthetic evaluation, and satisfaction rates. Then a data extraction sheet was set in Excel (Microsoft, Redmond, Washington, USA). Additionally, each article was assessed for the risk of bias in accordance with the methodological standards listed in the non-comparative case series checklist (for case series) and Ottawa-Newcastle Scale (for cohort studies), respectively (<https://www.ncbi.nlm.nih.gov/books/NBK35156/>) (Table S1, Table S2).

enucleation in Ghanaian children: 5 years' experience. *Ghana Med J*. 2014;48(4):204-7. Epub 2015/02/25. doi: 10.4314/gmj.v48i4.6. PubMed PMID: 25709135; PubMed Central PMCID: PMC4335434. 19. Jason D. Meier MRAG, MD; Mark J. Glasgold, MD. Autologous Fat Grafting Long-term Evidence of Its Efficacy in Midfacial Rejuvenation. 20. Bernardini FP, Gennai A, Izzo L, Zambelli A, Repaci E, Baldelli I, et al. Superficial Enhanced Fluid Fat Injection (SEFFI) to Correct Volume Defects and Skin Aging of the Face and Periocular Region. *Aesthetic surgery journal*. 2015;35(5):504-15. Epub 2015/04/26. doi: 10.1093/asj/sjv001. PubMed PMID: 25911629. 21. Kim HS, Choi CW, Kim BR, Youn SW. Effectiveness of Transconjunctival Fat Removal and Resected Fat Grafting for Lower Eye Bag and Tear Trough Deformity. *JAMA Facial Plast Surg*. 2019;21(2):118-24. Epub 2018/11/13. doi: 10.1001/jamafacial.2018.1307. PubMed PMID: 30418468; PubMed Central PMCID: PMC6439811. 22. Huang SH, Lin YN, Lee SS, Huang YH, Takahashi H, Chou CK, et al. Three Simple Steps for Refining Transcutaneous Lower Blepharoplasty for Aging Eyelids: The Indispensability of Micro-Autologous Fat Transplantation. *Aesthet Surg J*. 2019;39(11):1163-77. Epub 2019/01/23. doi: 10.1093/asj/sjz005. PubMed PMID: 30668643. 23. Kim J, Shin H, Lee M, Shin D, Kim S, Jo D, et al. Percutaneous Autologous Fat Injection Following 2-Layer Flap Lower Blepharoplasty for the Correction of Tear Trough Deformity. *The Journal of craniofacial surgery*. 2018;29(5):1241-4. Epub 2018/04/03. doi: 10.1097/scs.0000000000004552. PubMed PMID: 29608475. 24. Chen H, Zhang Q, Qiu Q, Yang Z. Autologous Fat Graft for the Treatment of Sighted

Posttraumatic Enophthalmos and Sunken Upper Eyelid. Ophthalmic Plast Reconstr Surg. 2018;34(4):381-6. Epub 2018/01/26. doi: 10.1097/IOP.0000000000001028. PubMed PMID: 29369151. 25. Miranda SG, Codner MA. Micro Free Orbital Fat Grafts to the Tear Trough Deformity during Lower Blepharoplasty. Plast Reconstr Surg. 2017;139(6):1335-43. Epub 2017/02/16. doi: 10.1097/PRS.0000000000003356. PubMed PMID: 28198772. 26. Ma Z, Chen G, Huang J, Wang J, Huang W. Clinical application of orbital fat autograft for tear trough deformity through tranconjunctival approach. Chin J Plast Surg. 2017;33(2):116-9. 27. Gennai A, Zambelli A, Repaci E, Quarto R, Baldelli I, Fraternali G, et al. Skin Rejuvenation and Volume Enhancement with the Micro Superficial Enhanced Fluid Fat Injection (M-SEFFI) for Skin Aging of the Periocular and Perioral Regions. Aesthetic surgery journal. 2017;37(1):14-23. Epub 2016/06/01. doi: 10.1093/asj/sjw084. PubMed PMID: 27241362. 28. Chiu CY, Shen YC, Zhao QF, Hong FL, Xu JH. Treatment of Tear Trough Deformity: Fat Repositioning versus Autologous Fat Grafting. Aesthetic plastic surgery. 2017;41(1):73-80. Epub 2016/12/23. doi: 10.1007/s00266-016-0692-z. PubMed PMID: 28008460. 29. Jiang J, Wang X, Chen R, Xia X, Sun S, Hu K. Tear trough deformity: different types of anatomy and treatment options. Postepy dermatologii i alergologii. 2016;33(4):303-8. Epub 2016/09/09. doi: 10.5114/ada.2016.61607. PubMed PMID: 27605904; PubMed Central PMCID: PMC5004220. 30. Park S, Kim B, Shin Y. Correction of superior sulcus deformity with orbital fat anatomic repositioning and fat graft applied to retro-orbicularis oculi fat for Asian eyelids. Aesthetic plastic surgery

Aesthetic plastic surgery. 2011;35(2):162-70. Epub 2010/09/14. doi: 10.1007/s00266-010-9574-y. PubMed PMID: 20835821.

31. Litwin AS, Poitelea C, Tan P, Ziahosseini K, Malhotra R. Complications and outcomes of grafting of posterior orbital fat into the lower lid-cheek junction during orbital decompression. Orbit (Amsterdam, Netherlands). 2018;37(2):128-34. Epub 2017/10/13. doi: 10.1080/01676830.2017.1383452. PubMed PMID: 29023175.

32. Chang HS, Lee D, Taban M, Douglas RS, Goldberg RA. "En-glove" lysis of lower eyelid retractors with AlloDerm and dermis-fat grafts in lower eyelid retraction surgery. Ophthalmic Plast Reconstr Surg. 2011;27(2):137-41. Epub 2010/06/22. doi: 10.1097/IOP.0b013e3181c53d38. PubMed PMID: 20562664.

33. Çoban Karataş M, Yaycıoğlu RA, Canan H. Orbital Dermis-Fat Graft Transplantation: Results in Primary and Secondary Implantation. Türk Oftalmoloji Dergisi. 2015;45(2):65-70. doi: 10.4274/tjo.55823.

34. Roh MR, Kim TK, Chung KY. Treatment of infraorbital dark circles by autologous fat transplantation: a pilot study. The British journal of dermatology. 2009;160(5):1022-5. Epub 2009/05/13. doi: 10.1111/j.1365-2133.2009.09066.x. PubMed PMID: 19434788.

35. Youn S, Shin JI, Kim JD, Kim JT, Kim YH. Correction of infraorbital dark circles using collagenase-digested fat cell grafts. Dermatologic surgery : official publication for American Society for Dermatologic Surgery [et al]. 2013;39(5):766-72. Epub 2013/02/27. doi: 10.1111/dsu.12140. PubMed PMID: 23437990.

36. Pelle-Ceravolo M, Angelini M. Properly Diluted Fat (PDF): An Easy and Safe Approach to Periocular Fat Grafting. Aesthetic surgery journal. 2020;40(1):19-33. Epub 2019/02/13. doi: 10.1007/s00266-019-01111-1.

10.1093/asj/sjz039. PubMed
PMID: 30753276. 37. Park JY,
Kim N. Periorbital
Lipogranuloma after Facial
Autologous Fat Injection and
Its Treatment Outcomes.
Korean journal of
ophthalmology : KJO.
2016;30(1):10-6. Epub
2016/02/13. doi:
10.3341/kjo.2016.30.1.10.
PubMed PMID: 26865798;
PubMed Central PMCID:
PMCPMC4742640. 38. Li XQ,
Wang TL, Wang JQ. Ptosis: An
Underestimated Complication
after Autologous Fat Injection
into the Upper Eyelid.
Aesthetic surgery journal.
2015;35(6):Np147-53. Epub
2015/08/01. doi:
10.1093/asj/sjv015. PubMed
PMID: 26229134. 39. Rohrich
RJ, Mahedia M, Shah N, Afrooz
P, Vishvanath L, Gupta RK.
Role of Fractionated Fat in
Blending the Lid-Cheek
Junction. Plast Reconstr Surg.
2018;142(1):56-65. Epub
2018/06/08. doi:
10.1097/PRS.00000000000000
4526. PubMed PMID:
29878987. 40. Hsu VM,
Stransky CA, Bucky LP, Percec
I. Fat grafting's past, present,
and future: why adipose tissue
is emerging as a critical link to
the advancement of
regenerative medicine.
Aesthet Surg J.
2012;32(7):892-9. Epub
2012/09/04. doi:
10.1177/1090820X1245565
8. PubMed PMID: 22942117.
41. Behr B, Tang C, Germann
G, Longaker MT, Quarto N.
Locally applied vascular
endothelial growth factor A
increases the osteogenic
healing capacity of human
adipose-derived stem cells by
promoting osteogenic and
endothelial differentiation.
Stem Cells. 2011;29(2):286-
96. Epub 2011/07/07. doi:
10.1002/stem.581. PubMed
PMID: 21732486; PubMed
Central PMCID:
PMCPMC3400547. 42.
Tonnard P, Verpaele A,
Peeters G, Hamdi M,
Cornelissen M, Declercq H.
Nanofat grafting: basic
research and clinical
applications. Plastic and
reconstructive surgery.
2013;132(4):1017-26. Epub

2013/06/21. doi:
10.1097/PRS.0b013e31829f
e1b0. PubMed PMID:
23783059. 43. Jiang S, Quan
Y, Wang J, Cai J, Lu F. Fat
Grafting for Facial
Rejuvenation Using Stromal
Vascular Fraction Gel Injection.
Clin Plast Surg.
2020;47(1):73-9. Epub
2019/11/20. doi:
10.1016/j.cps.2019.09.001.
PubMed PMID: 31739900. 44.
Ho Quoc C, Taupin T, Guerin
N, Delay E. Volumetric
evaluation of fat resorption
after breast lipofilling. Ann Chir
Plast Esthet. 2015;60(6):495-
9. Epub 2015/08/01. doi:
10.1016/j.anplas.2015.06.01
1. PubMed PMID: 26229038.
45. Stig-Frederik Trojahn Kølle
AF-N, Anders Bruun
Mathiasen, Jens Jørgen Elberg,
Roberto S Oliveri, Peter V
Glovinski, Jens Kastrup, Maria
Kirchhoff , Bo Sonnich
Rasmussen, Maj-Lis Møller
Talman, Carsten Thomsen,
Ebbe Dickmeiss, Krzysztof
Tadeusz Drzewiecki.
Enrichment of autologous fat
grafts with ex-vivo expanded
adipose tissue-derived stem
cells for graft survival: a
randomised placebo-controlled
trial. Lancet. 2013;382:1113-
20. 46. Gir P, Brown SA, Oni
G, Kashefi N, Mojallal A,
Rohrich RJ. Fat grafting:
evidence-based review on
autologous fat harvesting,
processing, reinjection, and
storage. Plast Reconstr Surg.
2012;130(1):249-58. Epub
2012/06/30. doi:
10.1097/PRS.0b013e318254
b4d3. PubMed PMID:
22743888. 47. Pusic AL,
Klassen AF, Scott AM, Klok JA,
Cordeiro PG, Cano SJ.
Development of a new
patient-reported outcome
measure for breast surgery:
the BREAST-Q. Plast Reconstr
Surg. 2009;124(2):345-53.
Epub 2009/08/01. doi:
10.1097/PRS.0b013e3181ae
e807. PubMed PMID:
19644246.

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Feb 26, 2021

Last Modified: Feb 26, 2021

PROTOCOL integer ID:
47696

Keywords: Eyelid, Periorbital area, Fat grafting, Adipose-derived stem cell, Facial rejuvenation

- 1 We will search the PubMed, Cochrane library, and Embase databases for articles published prior to November 11, 2020.
- 2 We will scrutinize the titles, abstracts, and full text of the retrieved articles and screen out articles that do not meet the inclusion and exclusion criteria.
- 3 We will perform meta-analysis by a comprehensive meta-analysis software (CMA), version 2.2.050 (Biostat, Englewood, NJ, USA). A random-effects meta-analysis will be conducted to analyze all the evidence.