VERSION 2

APR 21, 2021

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bucnnsve

Protocol Citation: Michelle Newbery, Simon Maksour, Amy Hulme, Neville S Ng, Mirella Dottori, Lezanne Ooi 2021. Efficient third generation lentiviral particle production. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bucnnsve> Version created by Neville S Ng

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
Lentiviral particle production method optimised for cost, time, and yield, compatible with animal-free and serum-free cell culture and accessible to basic cell culture laboratories. Supports multiple freeze-thaw cycles of viral particles without loss of transduction efficiency.

Created: Apr 20, 2021

Efficient third generation lentiviral particle production V.2

Michelle Newbery^{1,2}, Simon Maksour^{1,3}, Amy Hulme^{1,3},
Neville S Ng^{1,2}, Mirella Dottori^{1,3,4}, Lezanne Ooi^{1,2,4}

¹Illawarra Health and Medical Research Institute, Wollongong, NSW, Australia;

²School of Chemistry and Molecular Bioscience, University of Wollongong, Wollongong, NSW, Australia;

³School of Medicine, University of Wollongong, Wollongong, NSW, Australia;

⁴Molecular Horizons, University of Wollongong, Wollongong, NSW, Australia

Michelle Newbery: Authors contributed equally;

Neville S Ng: Authors contributed equally;

Lezanne Ooi: Corresponding author;

Neville S Ng

ABSTRACT

The overexpression of a gene of interest by third generation lentiviral particle generation systems is a critical process in molecular biology, cell biology and gene therapy research. While many lentiviral protocol production methods have been discussed in literature, this protocol takes into account previously established optimisations with the aim to minimise user handling time, cost, and maximise practical yield. This protocol allows for at least 6 days of consecutive viral particle collection without compromising HEK293T cell culture or viral production efficiency, and can be easily and cost effectively reproduced in basic cell culture laboratories.

PROTOCOL integer ID:
49262

Keywords: lentiviral,
HEK293T, differentiation,
lentivirus, overexpression,
plasmid, viral particle,
production

Cell culture

HEK293T cell culture

FreeStyle™ 293 Expression Medium (Thermo Fisher Scientific #12338018) OR
DMEM/F12 (Thermo Fisher Scientific #21331020) + GlutaMAX™ Supplement
(Thermo Fisher Scientific #35050061) + generic fetal calf serum OR Gibco KnockOut™
Serum Replacement (Thermo Fisher Scientific #10828028)
Gibco Penicillin-Streptomycin (Thermo Fisher Scientific #15140122)

Transfection

Linear 10-20 kDa PEI (Sigma #765090 or Polysciences #23966-1)
Plasmid(s) containing gene of interest

Plasticware

Generic tissue culture 75 cm² cell culture vessels and 10 mL serological pipettes
Generic round bottom ultracentrifuge tubes ~35 mL
Generic sterile 50 mL centrifuge tubes
Generic 50 mL syringe and 0.45 µm pore PES filters

SAFETY WARNINGS

⚠ All stages should be performed with appropriate safety precautions specific to local standards. Lentivirus particles can be inactivated by hypochlorite, peroxide and ethanol based sterilisation agents.

Lentiviral transfection

- 1 Maintain HEK293T cells in 0.15 mL/cm² (e.g. 12 mL in 75 cm² flask) animal product free FreeStyle 293 Expression Medium, foetal serum free DMEM/F12 + 2 mM GlutaMAX + 5% Knock Out Serum Replacement or DMEM/F12 + 2 mM GlutaMAX + 5% FBS. Include 50 U/mL Penicillin and Streptomycin to reduce risk of bacterial contamination if necessary.

Dissociate and subculture HEK293T to ~70% confluency per virus of interest.

Viral particle production can be performed within at least 15 passages without loss of yield.

CITATION

Ausubel LJ, Hall C, Sharma A, Shakeley R, Lopez P, Quezada V, Couture S, Laderman K, McMahon R, Huang P, Hsu D, Couture L (2012). Production of CGMP-Grade Lentiviral Vectors.. BioProcess international.

- 2 For each lentiviral plasmid containing gene of interest, calculate reagent volumes required for preparation of 2.5:1 PEI:DNA, 60 µg PEI per 75 cm².

Maximal transfection efficiency by PEI complex is typically obtained with higher proportions of PEI:DNA (Figure 1). We observed no differences in transfection in KSR or FBS -based medium.

Figure 1. Approximately 2.5-5:1 PEI:DNA provides maximal transfection efficiency. Experiment performed with a hNGN2-eGFP containing plasmid (n = 3, error presented as SEM; ** p < 0.01, *** p < 0.001, analysis by 1-way ANOVA and Holm-Sidak post-hoc multiple comparisons test).

- 3 Prepare PEI-DNA solution in 1.5 mL of DMEM/F12 and incubate for 00:05:00 at ambient temperature. 5m

- 4 Add PEI-DNA complex solution to cell culture flask(s) and return to incubator Overnight .

Note PEI-DNA complex transfection can occur without usage of low serum transfection medium products.

PEI-DNA complex precipitate should be microscopically visible as ~0.3 µm objects (Figure 2).

Figure 2. Phase contrast image of HEK293T cell culture with PEI:DNA complex particles.

CITATION

González-Domínguez I, Grimaldi N, Cervera L, Ventosa N, Gòdia F (2019). Impact of physicochemical properties of DNA/PEI complexes on transient transfection of mammalian cells.. New biotechnology.

LINK

<https://doi.org/10.1016/j.nbt.2018.09.005>

- 5** Each day collect medium in an appropriate sized sterile container (e.g. 50 mL centrifuge tubes or 120 mL specimen collection tubes), and store at 4 °C . Proceed to viral particle concentration at selected endpoint (Step 6).

We utilise viral particle concentration to avoid prolonged incubation of pluripotent or multipotent stem cells in HEK293T culture medium during viral transduction experiments. If viral particle concentration is not necessary, freeze immediately at -80 °C .

Note a single collection step is performed on day 6 due to negligible difference in loss of yield between day 3 and 6, which avoids multiple rounds of centrifugation.

CITATION

Rahman, H.; Taylor, Josephine; Clack, Beatrice; Stewart, R. S.; and Canterbury, Sarah C (2013). Effects of Storage Conditions on the Morphology and Titer of Lentiviral Vectors. SFA ScholarWorks.

Viral particle collection can be performed for at least up to 7 days.

CITATION

Ichim CV, Wells RA (2011). Generation of high-titer viral preparations by concentration using successive rounds of ultracentrifugation.. Journal of translational medicine.

LINK

<https://doi.org/10.1186/1479-5876-9-137>

Lentiviral particle concentration

1h 33m

- 6** Sterilise ultracentrifuge tubes and appropriate sized viral particle aliquot tubes (0.2 or 1.5 mL microcentrifuge tubes).

For each 75 cm² flask if collecting 12 mL over 6 days, at least 2 x 36 mL ultracentrifuge tubes are

required.

- 7 Centrifuge pooled viral particle supernatant at 300 x g, 00:03:00 to remove cell debris. 3m
- 8 Filter supernatant from each tube through a 0.45 µm pore PES filter into corresponding ultracentrifuge tubes.
- 9 Balance ultracentrifuge tubes within 0.1 g by weighing and adding appropriate amount of medium.
- 10 Collect viral particles by centrifugation of filtered viral particle supernatant at 50000 x g, 4°C, 01:30:00. 1h 30m
- 11 Carefully transfer tubes on ice, mark pellet position/expected position of pellet and decant slowly to discard supernatant.
- 12 Resuspend pellet as a ~200X concentrate of total supernatant (e.g. 180 µL from 36 mL of supernatant).

As lentivirus particles are enveloped and more susceptible to degradation in comparison to unenveloped DNA viruses, the addition of 10% DMSO as a cryoprotectant allows at least 4X freeze-thaw cycles without compromising viral particles and reduces necessity for low volume aliquots (Figure 3).

Figure 3. Inclusion of DMSO retains transfection efficiency of viral particle aliquots following 4X freeze-thaw cycles. Experiment performed with a hNGN2-eGFP containing plasmid (n = 3, error presented as SEM; * p < 0.05, comparison with Student's t-test).

13 Prepare viral particle aliquots of appropriate volumes corresponding to application and freeze at

-80 °C .

Viral titre (TU/mL) can be calculated (# transfected cells / volume of virus) based on fluorescent reporter or immunofluorescent staining.