APR 24, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.2angade

Protocol Citation: Lawrence A. Klobutcher, Larry Klobutcher 2019. Electroporation of *Euplotes crassus*: conditions that allow for the uptake of DAPI.. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.2angade>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Apr 24, 2019

Last Modified: Apr 24, 2019

PROTOCOL integer ID:
 22574

Electroporation of *Euplotes crassus*: conditions that allow for the uptake of DAPI.

Forked from [Electroporation of *Euplotes crassus*: conditions that allow for the uptake of DAPI.](#)

In 1 collection

Lawrence A. Klobutcher¹, Larry Klobutcher²

¹University of Connecticut Health Center; ²University of Connecticut, School of medicine

Protist Research to Optimize Tools in Genetics (PROT-G)

Angela Piersanti
 University of Camerino

ABSTRACT

The electroporation procedure and conditions described allow for the entry of DAPI into the cell, resulting in nuclear staining. We have not yet been successful in using these conditions for transformation of *Euplotes crassus*, but only a couple of constructs have been tried to date.

1 Grow a 600 ml culture of *Euplotes crassus* using [Dunaliella salina grown in artificial seawater](#) as the food source until all the algae is eaten, and then allow the *Euplotes* culture to starve an additional 24 hours.

2 Collect and concentrate cells down to ~40 ml by pouring the culture through a 20 um Nitex membrane.

Photo of 20um Nitex membrane and a frame constructed by cutting out the top and bottom of a Tupperware container.

- 3 Transfer the concentrated *Euplotes* cells to a 50 ml screw cap centrifuge tube and pellet cells by centrifugation at 1,000 rpm ($\sim 200 \times g$) for 1.5 min in a clinical centrifuge.
- 4 Remove all but the last 10 ml of the supernatant, resuspend cell pellet by gently flicking tube, and transfer resuspended cells to a 15 ml centrifuge tube.
- 5 Centrifuge cells at $200 \times g$ for 1.5 min, remove all but the last ~ 100 μ l of supernatant, and resuspend pelleted cells by gently agitating tube.
- 6 Add 1 ml of freshly prepared 0.3 M glucose to the tube of cells and gently mix.
- 7 Allow cells to equilibrate for 15 min at room temperature.
- 8 For electroporation, transfer 300 μ l of cells to a 0.2 cm electroporation cuvette.
- 9 Add 3 μ l of a 10 mg/ml solution of DAPI (or DNA) to the cells and gently agitate to mix.

- 10 Place cuvette in BTX ECM 630 electroporation device and electroporate using either 150 volts, 25 ohms, and 50 uF or 250 volts, 25 ohms, and 50 uF. Allow cells to recover for at least one min.
- 11 Transfer electroporated cells to a 15 ml screw-cap tube containing either 10 ml of a *Dunaliella* algae culture grown in artificial seawater to a low-moderate density or 9 ml of 0.3 M glucose + 1 ml of a moderate-high density algae culture.
- 12 Examine a sample of cells using a fluorescent microscope equipped with a DAPI filter. About 50% of cells are killed using these electroporation conditions, but a high percentage of the remaining cells should be motile and have brightly staining C-shaped macronuclei. Control, non-electroporated cells will have very dim macronuclear staining, with the exception of any dead cells present in the culture.

- 13 NOTES: The electroporation procedure and conditions described allow for the entry of DAPI into the cell, resulting in nuclear staining. We have not yet been successful in using these conditions for transformation of *Euplotes crassus*, but only a couple of constructs have been tried to date.
In step 11, the 0.3 M glucose + algae solution should be used if subsequent selection of transformants with a drug will be attempted, as we have found that this increases the sensitivity of *E. crassus* to at least some selective agents (e.g., G418 or paromomycin).