VERSION 1

APR 07, 2019

Electrotransformation of Clostridium species V.1

Chin Yee Tan¹

¹Duke University

Chin Yee Tan

ABSTRACT

A brief protocol for electrotransformation of Clostridium species

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.zvcf62w

Protocol Citation: Chin Yee Tan 2019.

Electrotransformation of Clostridium species.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.zvcf62w>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: In development
We are still developing and optimizing this protocol

Created: Apr 07, 2019

Last Modified: Apr 07, 2019

PROTOCOL integer ID:
22148

- 1 Inoculate 10 μL of BHI-HKC broth with 0.1 mL of stock culture overnight
- 2 Use the overnight culture to inoculate 100 mL of TPG to a starting density of OD 0.02
- 3 Harvest early-exponential phase culture (OD 0.2 to 0.25) by centrifugation by 12,000g x 00:15:00 at 25 °C
- 4 Wash once in 10 mL of SMP electroporation buffer
- 5 Resuspend in 10 mL of SMP
- 6 Take 0.4 mL of aliquots and mix with 500 ng of DNA, transfer to prechilled cuvettes with 0.2cm gap
- 7 Incubate on ice for 00:10:00
- 8 Electroporate at the following parameters: 25 μF , resistance 200 Ω , voltage 1.8kV

9 Immediately transfer the cells into 10 mL of BHI broth and incubate for 03:00:00 .

10 Plate cells in dilutions on solid selective and non-selective BHI agar