AUG 06, 2018

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.seeebbe

External link: <http://www.qa-bio.com/docs/E-EF02.QA-Bio.specsheet.pdf>

Protocol Citation: Mike Gibson 2018. Endo F2.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.seeebbe>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Aug 06, 2018

Last Modified: Aug 06, 2018

PROTOCOL integer ID:
 14502

Keywords: endoglycosidase
 f2, endo f2

Endo F2

Mike Gibson¹

¹QA-Bio Inc

Mike Gibson

ABSTRACT

Endo F2 cleaves Asparagine-linked high mannose or biantennary oligosaccharides. It cleaves between the two N-acetylglucosamine residues in the diacetylchitobiose core of the oligosaccharide, generating a truncated sugar molecule with one N-acetylglucosamine residue remaining on the asparagine. In contrast, PNGase F removes the oligosaccharide intact.

MATERIALS

MATERIALS

Endo F2 QA-Bio Inc Catalog #E-EF02

- 1** Add up to 200 µg of glycoprotein to an Eppendorf tube. Adjust to 38 µl final volume with de-ionized water.
- 2** Add 10 µl 5x Reaction Buffer 4.5
- 3** Add 2.0 µl of Endo F2 to the reaction. Incubate 3 hours at 37°C.
- 4** Monitor cleavage by SDS-PAGE.