



Jun 12, 2024

Version 2

# Enterovirus coxsackievirus A16 2A protease small scale expression and purification protocol V.2



Version 1 is forked from [SARS-CoV-2 Mpro small scale expression and purification protocol](#)

DOI

[dx.doi.org/10.17504/protocols.io.rm7vzj3y5lx1/v2](https://dx.doi.org/10.17504/protocols.io.rm7vzj3y5lx1/v2)

Korvus Wang<sup>1</sup>, Michael Fairhead<sup>1</sup>, Eleanor Williams<sup>1</sup>

<sup>1</sup>Centre for Medicines Discovery, University of Oxford

ASAP Discovery



Korvus Wang

CMD

## Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.rm7vzj3y5lx1/v2>

**Protocol Citation:** Korvus Wang, Michael Fairhead, Eleanor Williams 2024. Enterovirus coxsackievirus A16 2A protease small scale expression and purification protocol. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.rm7vzj3y5lx1/v2> Version created by **Korvus Wang**

**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working

**We use this protocol and it's working**

**Created:** June 12, 2024

**Last Modified:** June 12, 2024

**Protocol Integer ID:** 101676

**Keywords:** expression, purification, ASAP, AViDD, CMD, Enterovirus, coxsackievirus, coxsackievirus A16, 2A protease, purification of coxsackievirus a16, enterovirus coxsackievirus a16, picornaviridae enterovirus coxsackievirus a16, coxsackievirus a16, 2a protease, purification protocol this protocol, cva16, purification protocol, purification, protocol details the expression, sumo tag, sumo tag at small scale

**Funders Acknowledgements:**

National Institutes of Health/National Institute Of Allergy and Infectious Diseases (NIH/NIAID)

Grant ID: U19AI171432

## Disclaimer

Research was supported in part by NIAID of the U.S National Institutes of Health under award number U19AI171399. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

## Abstract

This protocol details the expression and purification of coxsackievirus A16 2A protease construct bearing a N-terminal His-SUMO tag at small scale (<6L).

Picornaviridae Enterovirus coxsackievirus A16 (CVA16)

## Attachments



PAGE23-01675 - Enter...

281KB

## Guidelines

- **Construct / plasmid resource-name:** A71EV2A protease construct bearing a N-terminal SUMO-His tag.
- **Construct Addgene ID:** Processing

## Materials

### Plasmid details:

Addgene plasmid processing

- Vector: pNIC
- Cell line: *E. coli* Rosetta strain BL21(DE3)-RR
- Tags and additions: N-terminal His-SUMO tag
- Construct protein sequence:  
 MHHHHHHGSGDQEAKPSTEDLGDKKEGEYIKLKVIGQDSSEIHFVKMTTHLKKLKESYCQRQGVPMNSLRFLFEGQRI  
 ADNHTPKELGMEEDVIEVYQEQTGGSGAIYVGNYRVVNRHLATHNDWANLVWEDSSRDLLVSSTTAQGCDTIARCDC  
 QTGVYYCSSRRKHYPVSFSKPSLIFVEASEYYPARYQSHLMLAVGHSEPGDCGILRCQHGTVGVIVSTGGNGLVGFADVR  
 DLLWLDEEAMEQ

### Expression

**AIM-TB:** TB autoinduction media (Formedium AIMTB0210, ordered without added glucose and lactose)

After autoclaving, add 20mL of 50x AIM mix (400mL glycerol, 100g lactose, 25g glucose in 1L of ddH<sub>2</sub>O, filter sterilised) per L of media

### Purification

Chicken hen egg white lysozyme (Merck, 62971)

Benzonase (Merck, 1.01654)

Imidazole (Merck, RDD044)

Ni Sepharose 6 FF resin (Cytiva, 17531801)

Gravity flow column, 2.5cm diameter (Bio Rad, 7372532)

Centrifugal concentrators, 10kDa MWCO (Merck, UFC901008)

On an FPLC system:

On an FPLC system:

SEPAX SEC SRT-100 (Sepax Tech, 215100-21230)

or

HiLoad 16/600 Superdex 75 pg (Cytiva, 28989333)

5mL sample loop

SDS-PAGE sample buffer, gel, and gel tank

### Lysis buffer:

	A	B
	Hepes (pH 7.5)	50 mM



	A	B
	NaCl	500 mM
	Glycerol	5%
	TCEP	0.5 mM
	Lysozyme	0.5 mg/mL
	Benzonase	0.05 mg/mL

Prepare 100 mL per 1 L *E.coli* expression

**Base buffer:**

	A	B
	Hepes (pH 7.5)	50 mM
	NaCl	50 mM
	Glycerol	5%
	TCEP	0.5 mM

Prepare 2 L per 6 L *E.coli* expression. Used to prepare the following buffers

**Binding buffer:** base buffer

**Wash buffer 1:** base buffer + 30mM imidazole

**Wash buffer 2:** base buffer + 50mM imidazole

**Elution buffer:** base buffer, add 500mM imidazole

**Gel filtration buffer:**

	A	B
	Hepes (pH 7.5)	25 mM
	NaCl	300 mM
	Glycerol	5%
	TCEP	0.5 mM

**SDS-PAGE:** NuPage 4-12%, Bis-Tris protein gel, 26 well (Thermo-Fisher, WG1403BOX)

Run in MES buffer, 200V 35mins.

## Troubleshooting



## Abbreviations

- 1 CV - column volume, total volume of resin in a column  
IMAC - immobilised metal affinity chromatography  
FT - flow through

## Plasmid Transformation

1d

- 2 Transform the coxsackievirus A16 2A protease plasmid into BL21(DE3) and store a glycerol stock of this at -80 °C  
The coxsackievirus A16 2A protease plasmid encodes the 2A protease with an N-terminal His6-SUMO tag on a kanamycin resistant backbone with a T7 promoter.

## Protein expression

2d 10h

- 3 Scrape off some of the glycerol stock with a sterile loop and use this to inoculate a 50 mL falcon tube containing 10 mL of LB supplemented with 50 Mass Percent kanamycin. Grow the starter culture at 37 °C Overnight with 200 rpm shaking.
- 4 Use the 10 mL starter culture to inoculate 1 L Sample (see Materials) supplemented with 50 Mass Percent kanamycin in a baffled flask.  
 200 rpm, 37°C

1d

6h



### Note

For this protocol typically 6 L of culture is grown for each purification

### Note

Media and expression condition are especially important for the production of this construct, as it is prone to auto-cleavage of the affinity tag during expression.



- 5 When the OD<sub>600</sub> reaches approximately 3.0, lower the temperature and shaker speed to 180 rpm, 18°C and incubate Overnight 1d
- 6 Harvest the cells by centrifugation at 4000 x g, 4°C, 00:30:00 . Discard the supernatant and store the pellet at -80 °C . 30m

#### Note

For reference: total pellet weight from 6L TB media should be around 90g.

## Protein Purification 2d

### 7 Lyse cell pellet 2h 30m

#### 7.1 1h

#### Note

See Materials tab for buffer compositions.

#### Note

##### Coxsackievirus A16 2A protease His6-SUMO fusion protein properties

Before tag cleavage:

MW=27.927 kDa

E (assume all Cys reduced)=32890 mM<sup>-1</sup>cm<sup>-1</sup>

PI=5.81

After tag cleavage:

MW=15.876 kDa

E (assume all Cys reduced)=28975 mM<sup>-1</sup>cm<sup>-1</sup>

PI=5.58

These values are determined by ExPASy ProtParam



Thaw and resuspend the pellet using ~7mL of lysis buffer per g of pellet. Stir gently at Room temperature for 00:30:00 to allow lysozyme and benzonase to start breaking down cell components.

7.2 Lyse cells by sonication on ice 00:00:04 On 00:00:12 Off for a total 'on' time of 00:07:00 at 50% amplitude to fully rupture the cells. Ensure pellet remains cold during sonication to prevent overheating.

7m 16s

7.3 Centrifuge the lysed cells for 38000 x g, 4°C, 01:00:00 to remove insoluble cell debris, and collect the soluble fraction in a bottle 4 °C

1h

## 8 Perform IMAC to extract target protein from the lysed cell mixture

8.1 Dispense 5 mL of IMAC resin (Ni Sepharose 6 FF, Cytiva) into a gravity flow column. Rinse resin with ~ 10 µL distilled water to remove the storage solution and then ~ 10 µL binding buffer to equilibrate the resin.

10m

8.2 Resuspend the equilibrated resin with 10 mL of binding buffer and add to the supernatant bottle. Incubate the resin with the soluble fraction for 00:30:00 while rotating or otherwise mixing gently at 4 °C

30m

8.3 Load the resin/soluble fraction mix back onto the gravity flow column, retaining the flow through separately for SDS-PAGE analysis.

30m

### Note

For SDS-PAGE samples, mix 15 uL sample with 5 uL 4x sample buffer, supplemented with 10 mM DTT.

8.4 Wash the column with 10 µL of base buffer, followed by 10 µL of wash buffer 1 and 2. Allow wash buffer to pass through completely between washes. This is to remove non-specific, weak binding of contaminant proteins from the resin for a cleaner elution. Collect washes separately for SDS-PAGE analysis.

30m

8.5 Elute the protein with 1.5 µL of elution buffer.


20m





8.6 Repeat step 8.5 one more time, collecting a total of 2 separate elution fractions. This is to ensure maximum retrieval of protein from the resin. The total protein concentration of the elutions are then measured by Nanodrop.

20m

8.7 Wash used IMAC resin with  10  $\mu\text{L}$  of base buffer, and leave the column submerged in a small amount of base buffer so that the resin is kept moist. This washed IMAC resin will later be reused for reverse IMAC (rIMAC)

9 Run SDS-PAGE of all samples from total lysis supernatant to final elution. Stain gel with protein staining solution Coomassie Blue and determine which fractions contain the target protein by finding the band corresponding to the target molecular weight.

40m

#### Note

The target protein is expected to be present mostly in the elution samples, although small amounts may be found in the FT and washes. If that is not the case, then further troubleshooting is required.

### 10 Elution de-salting, tag cleavage and reverse IMAC



1d

10.1 Pool the elutions and desalt using a HiPrep 26/10 desalting column, run on an AKTA pure at a maximum flow rate of 10mL/min.

30m

#### Note

Desalting reduces the concentration of imidazole in the sample which may inhibit SENP1 protease activity during tag cleavage as well as interfering with the reverse IMAC step.

10.2 Add His-SENP1 SUMO protease at a 1:100 ratio to the total protein content of the desalted sample, as determined by nanodrop. Incubate at  4 °C  Overnight This cleaves the affinity tag.


1d

10.3 Pour the cleaved Cocksackievirus A16 2A protease, SUMO tag, SENP1 protease mixture over the washed IMAC resin and collect the flow through, rIMAC.


30m

**Note**

This step will remove the cleaved tag and any un-cleaved target from the sample. As the SENP1 protease used is His-tagged, this is removed from the sample too.

- 10.4 Wash the IMAC resin with  2  $\mu$ L wash buffer 1 and 2 to remove any target protein still bound to the resin. Take samples of the FT and washes for SDS-PAGE analysis.


30m

- 10.5 (Optional) elute rIMAC resin with  2  $\mu$ L elution buffer to confirm if the protein shows non-specific binding to the resin used.

5m

**Note**

This will help determine if the protein is "sticky" to the IMAC resin matrix material, and help in further troubleshooting if the final yield is lower than expected. Despite having a His6 on it the cleaved SUMO tag is usually found contaminating the rIMAC fractions containing coxsackievirus A16 2A protease. The tag can however be removed in the subsequent gel filtration step.



**SDS-PAGE analysis of IMAC and cleavage fractions.** The higher molecular weight band in the right-hand-side gel image, highlighted by red arrow, agrees with the size of cleaved target protein, while the lower band highlighted with blue arrow corresponds to the size of the cleaved SUMO tag (12.372 kDa but band appears at ~15kDa on the SDS-PAGE gels used). Both bands are also observed throughout the IMAC fractions. This is because the coxsackievirus A16 2A protease construct used exhibits auto-cleavage activity during expression and purification.

## 11 Purify sample further by size exclusion chromatography.

6h

- 11.1 Using 10,000 MWCO spin concentrators, concentrate the rIMAC step containing fractions of the target protein to a final volume of under 5 mL .

1h

- 11.2 Remove any solid aggregates from the sample by centrifugation at 17200 x g, 4°C, 00:10:00 , then immediatly draw up the supernatant with a 5mL syringe and a blunt-tip fill needle, taking care not to disturb the pellet.

15m

### Note

This is to remove as much solid particles from the injection sample as possible, so as to not clog the in-line filter or frit of the column.


## 12 Using an AKTA Pure system:

2h


Inject the sample onto a 5mL sample loop and run the sample down HiLoad 16/60 Superdex 75 pg gel filtration column at 1 mL/min using gel filtration buffer as the mobile phase, collect 1mL fractions.

## 13 Analyze the size exclusion chromatography fractions by SDS-PAGE and pool the fractions with highest amounts of pure 2A protease.

1h



**Chromatogram of coxsackievirus A16 2A protease SEC run:** Fractions F9-H8 were analyzed by SDS-PAGE to see which contained the target protein




**SDS-PAGE analysis of SEC fractions F9-H8.** Fractions G8-H8 were pooled as they contain majority coxsackievirus A16 2A protease in comparison to contaminating SUMO.

- 13.1 Take the fractions that contain the cleanest target protein and concentrate to [M] 15 mg/mL using a 10 kDa MWCO centrifugal concentrator

10m

Take 1  $\mu$ L of the final sample for SDS-PAGE, and another for mass spectroscopy (MS).



**Intact mass spectroscopy result of the purified coxsackievirus A16 2A protease sample.** No mass corresponding to the SUMO tag (12.05 kDa) is observed in the final sample, only that of the untagged coxsackievirus 2A protease, 15.88 kDa.

- 13.2 Aliquot into appropriate volumes for future usage to minimise freeze/thaw cycles. Flash-freeze in liquid nitrogen, and store at  $-80^{\circ}\text{C}$  until required.

30m

For example:

The final yield from processing 6 L of cells was 50 mg of pure coxsackievirus A16 2A protease

