FEB 06, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.babriam6

External link:
<https://doi.org/10.1371/journal.pone.0227826>

Protocol Citation: Alison Lovegrove 2020. Enzymatic fingerprinting of AX and mixed-linked BG. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.babriam6>

MANUSCRIPT CITATION:
 Lovegrove A, Wingen LU, Plummer A, Wood A, Passmore D, Kosik O, Freeman J, Mitchell RAC, Hassall K, Ulker M, Tremmel-Bede K, Rakszegi M, Bedő Z, Perretant M, Charmet G, Pont C, Salse J, Waite ML, Orford S, Burridge A, Pellny TK, Shewry PR, Griffiths S (2020) Identification of a major QTL and associated molecular marker for high arabinoxylan fibre in white wheat flour. PLoS ONE 15(2): e0227826. doi: [10.1371/journal.pone.0227826](https://doi.org/10.1371/journal.pone.0227826)

Enzymatic fingerprinting of AX and mixed-linked BG

PLOS One

Alison Lovegrove¹

¹Rothamsted Research

Alison Lovegrove

ABSTRACT

Lab protocol for: Enzymatic fingerprinting for arabinoxylan and mixed-linked beta-glucan in wheat flour by HPAEC-PAD. Based on Ordaz-Ortiz and Saulnier, 2005 J. Cereal Sci. 42. 119-125. and Odaz-Ortiz et al., 2005 J. Ag Food Chem. 53. 8349-8356.

ATTACHMENTS

[SOP. Fingerprinting of AX and MLG in wheat flour.docx](#)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: Dec 10, 2019

Last Modified: Feb 06, 2020

PROTOCOL integer ID:
30801