AUG 28, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bkexktfn

Protocol Citation: Angel A Justiz-Vaillant 2020. Enzyme linked immunosorbent assay (ELISA) for determining the plasma or serum concentration of IL-33 in humans.. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bkexktfn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Aug 28, 2020

Last Modified: Aug 28, 2020

PROTOCOL integer ID:
 41143

🌐 Enzyme linked immunosorbent assay (ELISA) for determining the plasma or serum concentration of IL-33 in humans.

Angel A Justiz-Vaillant¹

¹University of the West Indies St. Augustine

University of the West Indies

angel.vaillant@sta.uwi.edu

Angel A Justiz-Vaillant

University of the West Indies St. Augustine

DISCLAIMER

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](#) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](#), can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

- 1 Ninety-six well ELISA plates are coated with monoclonal anti-human antibodies to interleukin-33 (IL-33).

- 2 Patient serum samples are added to the plates.
- 3 The plate is incubate x 1.30 hour at RT.
- 4 The plate is washed 4 times with PBS-tween buffer.
- 5 The wells are incubated with a biotin conjugated anti-human IL-33 for 1.30 hour at RT.
- 6 The plate is washed again as above.
- 7 Add to the plate a peroxidase-labeled streptavidin conjugate and incubate it for 1 hour at RT.
- 8 After a further washing procedure a substrate solution reactive is added and allowed to produced a colored reaction in positive controls.
- 9 The level of IL-33 in the sample is proportional to the colored product developed.

- 10 The addition of 3M H₂SO₄ stops the reaction.
- 11 The absorbance is measured at 450 nm.
- 12 The IL-33 concentration can be calculated by generating an standard curve using lineal regression.