APR 24, 2019

🌐 Euplotes crassus transfection using FuGene HD Transfection Reagent as vehicle (provisional)

🔗 Forked from [Euplotes crassus transfection using FuGene HD Transfection Reagent as vehicle \(provisional\)](#)

📁 In 1 collection

RACHELE CESARONI¹, Rachele Cesaroni²

¹University of Bern, Institute of Biology; ²Universität Bern

Protist Research to Optimize Tools in Genetics (PROT-G)

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.2apgadn

Angela Piersanti
University of Camerino

Protocol Citation: RACHELE CESARONI, Rachele Cesaroni 2019. Euplotes crassus transfection using FuGene HD Transfection Reagent as vehicle (provisional).

protocols.io
<https://dx.doi.org/10.17504/protocols.io.2apgadn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: In development
We are still developing and optimizing this protocol

Created: Apr 24, 2019

Last Modified: Apr 24, 2019

PROTOCOL integer ID:
22575

- 1 Collect 4×10^4 well-fed Euplotes crassus cells (we used E.coli as the only food source) by centrifugation at 400 rcf for 3 minutes.

- 2 Wash the cells twice with artificial sea water (see attachment for the recipe) and once with 500 mM sorbitol, 0.5 mM Tris-HCl, pH 7.0 (400 rcf for 3 minutes each time). Then resuspend *Euplotes crassus* cells in 50 µl of the medium (500 mM sorbitol, 0.5 mM Tris-HCl, pH 7.0). To get this small volume you may require an additional minute of centrifugation.

Note

Recipe for complete sea water (1 L):

36 g Reef Crystals

1 ml Walne's solution

1 ml of 10 µg/ml FeSO₄

0.2 ml of 2 mg/ml thiamine (light sensitive; store at 4 °C)

Add distilled water up to 1 L

- 3 Transfer the cells into wells within a 96-well plate for transfection.
- 4 Allow FuGene HD Transfection Reagent to reach room temperature, and then mix it by inverting or vortexing briefly (if you see a precipitate, briefly warm it at 37°C, and then let it reach room temperature).
- 5 Add 90-98 µl of medium (500 mM sorbitol, 0.5 mM Tris-HCl, pH 7.0) at room temperature and 4 µg of DNA in MilliQ H₂O (0.2-1 µg/µL) to an Eppendorf tube and vortex (after adding the DNA the final volume must be 100 µl).
- 6 Add 6 µl of FuGene HD Transfection Reagent directly to the medium and mix immediately.
- 7 Incubate FuGene HD Transfection Reagent and DNA mixture for 15 minutes at room temperature.

- 8** Add 10 μl of the FuGene HD Transfection Reagent and DNA mixture to the 50 μl of *Euplotes crassus* cells in medium (mix everything by pipetting).
- 9** One hour after addition of FuGene/DNA complexes, add 50 μl of artificial sea water to the cells.
- 10** After another hour, add to the cells other 50 μl of artificial sea water.
- 11** An hour later harvest the cells (400 rcf for 3 minutes), and wash them twice with artificial sea water (400 rcf for 3 min each time). Then resuspend them in 400 μl of the artificial sea water.
- 12** Transfer the cells into glass depression wells for subsequent monitoring.
- 13** Incubate the cells at 24°C, then examine them by fluorescence microscopy to determine expression of the construct.