VERSION 2

MAY 19, 2021

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bu6anzae

Protocol Citation: Kazuma Rifu, Jun Watanabe, Hideki Sasanuma, Nobuyuki Taniguchi 2021. Evaluation of elasticity of the pancreas by Acoustic Radiation Force Impulse (ARFI) in acute pancreatitis patients: A systematic review and meta-analysis (protocol).
protocols.io
<https://dx.doi.org/10.17504/protocols.io.bu6anzae> Version created by Kazuma Rifu

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: May 19, 2021

Last Modified: May 19, 2021

Evaluation of elasticity of the pancreas by Acoustic Radiation Force Impulse (ARFI) in acute pancreatitis patients: A systematic review and meta-analysis (protocol) V.2

Kazuma Rifu¹, Jun Watanabe², Hideki Sasanuma³, Nobuyuki Taniguchi⁴

¹Department of Surgery, Division of Gastroenterological, General and Transplant Surgery, Jichi Medical University, Shimotsuke-City, Tochigi, Japan;

²Department of Surgery, Division of Gastroenterological, General and Transplant Surgery, Jichi Medical University, Shimotsuke-City, Tochigi, Japan. Division of Community and Family Medicine, Jichi Medical University, Shimotsuke-City, Tochigi, Japan.;

³Department of Surgery, Division of Gastroenterological, General and Transplant Surgery, Jichi Medical University, Shimotsuke-City, Tochigi, Japan.;

⁴Department of Clinical Laboratory Medicine, Jichi Medical University, 3311-1 Yakushiji, Shimotsuke, Tochigi, Japan.

Jichi Medical University

Kazuma Rifu
Jichi Medical University

ABSTRACT

Keywords: Acute
Pancreatitis, Acoustic
Radiation Force Impulse,
Elasticity Imaging Techniques

Acute pancreatitis (AP) is characterized by inflammation of the pancreas(1). The morbidity of AP is reported about 5-80/100000 people/year (2). An increase in the incidence of AP has been also reported (3-9). The mortality rate has been reported to be 1.7-8.4% (10, 11), and AP still remains a life-threatening disease. Clinical symptoms and elevated levels of pancreatic enzymes constitute the main diagnostic features of this condition.(12, 13) Early diagnosis and treatment are important, and reliable imaging techniques are required for a diagnosis of acute pancreatitis. Acoustic radiation force impulse (ARFI) elastography is a promising ultrasound-based imaging modality for the evaluation of tissue stiffness including in the diagnosis of AP (14). Studies using the ARFI elastography evaluated different abdominal organs(15), especially in pancreatic disease, such as AP, chronic pancreatitis, or pancreatic carcinoma, relevant higher shear wave velocities than in healthy pancreas parenchyma are described(1, 16, 17). In this study, we evaluated acoustic radiation force impulse (ARFI) elastography, Virtual Touch tissue quantification (VTQ; Siemens Medical Solutions, Mountain View, CA),to diagnose acute pancreatitis. The consistency of the pancreatic parenchyma usually becomes stiffer in acute pancreatitis as compared to the healthy pancreas, which is identifiable with Share Wave Elastography (SWE), including ARFI (18, 19). However, studies using elastographic techniques in patients with acute pancreatitis are conflicting(19).One prospective study failed to find significant differences in pancreatic shear wave velocities between patients with AP and healthy volunteers(14, 19). Three other studies showed significantly higher pancreatic shear wave velocities in patients with acute pancreatitis compared to persons with the normal pancreas(1, 18-20).

The present study will aim to assess the effectiveness of ARFI elastography in the form of ARFI-VTQ imaging in characterizing and differentiating normal pancreas and acute pancreatitis.

ATTACHMENTS

[KR\) ARFIprotocol
20210519.docx](#)