MAR 06, 2020

Extraction Procedure

PLOS One

Jing
Xu¹

¹Hainan University

Jing Xu
Hainan University

OPEN ACCESS

DOI:
[dx.doi.org/10.17504/protocols.io.bdawi2fe](https://doi.org/10.17504/protocols.io.bdawi2fe)

External link:
<https://doi.org/10.1371/journal.pone.0234435>

Protocol Citation: Jing Xu
2020. Extraction Procedure.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.bdawi2fe>

MANUSCRIPT CITATION:
Zhou J, Yang Q, Zhu X, Lin
T, Hao D, Xu J (2020)
Antioxidant activities of
Clerodendrum cyrtophyllum
Turcz leaf extracts and
their major components.
PLoS ONE 15(6): e0234435.
doi:
[10.1371/journal.pone.0234435](https://doi.org/10.1371/journal.pone.0234435)

License: This is an open
access protocol distributed
under the terms of
the [Creative Commons
Attribution License](#), which
permits unrestricted use,
distribution, and reproduction
in any medium, provided the
original author and source
are credited

Protocol status: Working
We use this protocol and it's
working

ABSTRACT

Dried leaves of *C. cyrtophyllum* (150 g) were weighted and sieved (20 mesh) in an herb disintegrator (118 Swing, Zhejiang, China); the powdered samples were extracted twice according to previous protocol [10]. Solvent was removed from the combined filtrates and 61.44 g of ECE was obtained and redissolved in distilled water (500 ml). The solution was partitioned with 3×250 ml petroleum ether (60–90 °C), 3×250 ml dichloromethane, 3×250 ml ethyl acetate and 3×250 ml *n*-butanol. The resulting extracts were concentrated to yield sub-fractions, PEF, DMF, EAF, BAF and RF, which were 0.65, 5.53, 4.13, 13.28 and 36.97 g, respectively. The samples were stored at 4 °C.

Created: Mar 06, 2020

Last Modified: Mar 06,
2020

PROTOCOL integer ID:
33846