OCT 14, 2019

Fatty acid extraction and derivatisation

Igem Dusseldorf¹

¹Heinrich-Heine Universität Düsseldorf

Igem Dusseldorf

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.767hrhn

Protocol Citation: Igem Dusseldorf 2019. Fatty acid extraction and derivatisation.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.767hrhn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: Oct 14, 2019

Last Modified: Oct 14, 2019

PROTOCOL integer ID:

28607

ABSTRACT

- C17 internal standard (stock = 1 mg/ml)
- MeOH//3N HCl
- hexane
- 1% NaCl
- clean glass tubes
- always work with glass pipettes or hamilton

Start

samples with 4 oD-units (e.g.: oD = 1, you will need 4 ml culture)
centrifuge in a clean glass tube for 10 min @ maximum speed
discard supernatant
Freeze @ -80 °C until further use

Extraction solution

200 µl C17 internal standard + 9,8 ml MeOH/3N HCl
caution! fill in a beaker and take the needed amount out of this beaker
always work with gloves
pipet C17 internal Std. with hamilton

Extraction

add 1 ml extraction solution to samples and blank
blank: fatty acid standard mix
heat 60 min @ 90°C
let cool down @ RT
add 1 ml hexane
add 1 ml 1% NaCl
vortex for 30 sec
spin down 5 min @ 2000 rpm
transfer hexane phase in to GC vial with screw cap
dilute samples (10 µl sample + 90 µl hexane)
wash hamilton between samples with hexane