



Sep 11, 2018 Version 2

## fecal bacterial DNA extraction V.2

 [PLOS One](#)

DOI

[dx.doi.org/10.17504/protocols.io.tfyejpw](https://dx.doi.org/10.17504/protocols.io.tfyejpw)

Shiu-Ming Kuo<sup>1</sup>

<sup>1</sup>University at Buffalo



Shiu-Ming Kuo

OPEN  ACCESS



DOI: [dx.doi.org/10.17504/protocols.io.tfyejpw](https://dx.doi.org/10.17504/protocols.io.tfyejpw)

External link: <https://doi.org/10.1371/journal.pone.0205055>

**Protocol Citation:** Shiu-Ming Kuo 2018. fecal bacterial DNA extraction. **protocols.io**

<https://dx.doi.org/10.17504/protocols.io.tfyejpw>

### Manuscript citation:

Zheng W, Wang K, Sun Y, Kuo S (2018) Dietary or supplemental fermentable fiber intake reduces the presence of *Clostridium XI* in mouse intestinal microbiota: The importance of higher fecal bacterial load and density. PLoS ONE 13(10): e0205055. doi: [10.1371/journal.pone.0205055](https://doi.org/10.1371/journal.pone.0205055)

**License:** This is an open access protocol distributed under the terms of the **[Creative Commons Attribution License](#)**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working

We use this protocol and it's working

**Created:** September 11, 2018

**Last Modified:** September 11, 2018

**Protocol Integer ID:** 15576

**Keywords:** feces; DNA; bacteria; microbiome

## Attachments



Fecal DNA Isolation ...

42KB

1