JUN 06, 2018

First-Strand cDNA Synthesis and Conventional Polymerase Chain Reaction (PCR)

Izabela M Rezende¹, Lívia Sacchetto

¹Stanford University School of Medicine

MRCA

Izabela M Rezende

Stanford University School of Medicine

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.pw9dph6

Protocol Citation: Izabela M Rezende, Lívia Sacchetto 2018. First-Strand cDNA Synthesis and Conventional Polymerase Chain Reaction (PCR). **protocols.io** <https://dx.doi.org/10.17504/protocols.io.pw9dph6>

ABSTRACT

For cDNA synthesis and for amplify larger fragments of YFV genome. The PCR target is of the CprM/envelope region of YFV genome described by Jorge and colleagues (2017).

MATERIALS

MATERIALS

- ☒ Microcentrifuge Tubes **Contributed by users**
- ☒ Filter Tips **Contributed by users**
- ☒ Thermocycler **Contributed by users**
- ☒ GoScript(TM) Reverse Transcriptase, 500 rxn **Promega Catalog #A5004**
- ☒ Random Primers, 20ug **Promega Catalog #C1181**
- ☒ Nuclease-Free Water, 1000ml **Promega Catalog #P1199**
- ☒ PCR tubes **Contributed by users**
- ☒ Primer **Contributed by users**
- ☒ Micropipettors **Contributed by users**
- ☒ GoTaq® Colorless Master Mix **Promega Catalog #M714**

**MANUSCRIPT CITATION:
Persistence of Yellow fever
virus outside the Amazon
Basin, causing epidemics in
Southeast Brazil, from
2016 to 2018.**

Rezende IM de*, Sacchetto
L,* Mello ÉM de, Alves
PA, Iani FC de M, Adelino
TÉR, Duarte MM, Cury, ALF,
Bernardes AFL, Santos TA,
Pereira LS, Dutra MRT,
Ramalho DB, Thoisy B, Kroon
EG, Trindade GS, Drumond BP
PLoS Neglected Tropical
Disease 2018;12:
e0006538. doi:10.1371/journ
al.pntd.0006538

*These authors contribully
equally to this work
Jorge, T. A.; Mosimann, A. L.
P.; Maron, L. N. A.; Santos, C.
N. D. dos (2017). Isolation
and characterization of a
Brazilian strain of yellow fever
virus from an epizootic
outbreak in 2009. Acta
Tropica, Volume
166, 2017, Pages 114-
120, ISSN 0001-
706X, [https://doi.org/10.1016/j](https://doi.org/10.1016/j.actatropica.2016.09.030)
.actatropica.2016.09.030.

GoScript™ Reverse
Transcription System
(Promega, USA), more
information available at:
[https://www.promega.com.br/](https://www.promega.com.br/-/media/files/resources/protocols/technical-manuals/101/goscript-reverse-transcription-system-protocol.pdf)
-
/media/files/resources/protocol
s/technical-
manuals/101/goscript-reverse-
transcription-system-
protocol.pdf

GoTaq® Colorless Master Mix
(Promega, USA), more
information available at:
[https://www.promega.com.br/](https://www.promega.com.br/-/media/files/resources/protocols/product-information-sheets/g/gotaq-colorless-master-mix.pdf)
-
/media/files/resources/protocol
s/product-information-
sheets/g/gotaq-colorless-
master-mix.pdf

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: May 05, 2018

Last Modified: Jun 06, 2018

PROTOCOL integer ID:
11969

First-Strand cDNA Synthesis

- 1 Close each tube of RNA tightly. Place tubes into a preheated 70°C heat block for 5 minutes. Immediately chill in ice-water for at least 5 minutes. Centrifuge each tube for 10 seconds in a microcentrifuge to collect the condensate and maintain the original volume. Keep the tubes closed and on ice until the reverse transcription reaction mix is added.
- 2 Mix and briefly centrifuge each component before use. Combine the following:

C o m p o n e n t	-
E x p e r i m e n t a l R N A	X μ L

Random Primer (0.5 μ g/reaction)	Nuclease-Free Water	Final volume
$\underline{X} \mu\text{L}$	$X \mu\text{L}$	$5 \mu\text{L}$

- 3** Prepare the reverse transcription reaction mix by combining the following components of the GoScript™ Reverse Transcription System in a sterile microcentrifuge tube on ice. Prepare sufficient mix to allow 15µl for each cDNA synthesis reaction to be performed. Determine the volumes needed for each component and combine them in the order listed. Vortex gently to mix, and keep on ice prior to dispensing into the reaction tubes.

Amount	Component
--------	-----------

Nuclease-Free Water (to a final volume of 15 μ l)	X μ l
Goscript [™] 5X Reaction Buffer	4.0 μ l

MgCl ₂ (final concentration 1.5 – 5.0 mM)	1.2 – 4.0 μ
PCR Nucleotide Mix (final concentration 0.5 mM each dNTP)	1.0 μ

GoScript™ Reverse Transcriptase	1.0 μ
---------------------------------------	----------

- 4 Add 15μl aliquots of the reverse transcription reaction mix to each reaction tube on ice. Be careful to prevent cross-contamination. Add 5μl of RNA and primer mix to each reaction for a final reaction volume of 20μl per tube.
- 5 Anneal: Place the tubes in a controlled-temperature heat block equilibrated at 25°C, and incubate for 5 minutes.
- 6 Extend: Incubate the tubes in a controlled-temperature heat block at 42°C for up to one hour. The extension temperature may be optimized between 37°C and 55°C. The reactions may be maintained frozen for long-term storage.
- 7 Inactivate Reverse Transcriptase: If the experimental goal is to proceed with PCR, the reverse transcriptase must be thermally inactivated prior to amplification. Incubate the reaction tubes in a controlled-temperature heat block at 70°C for 15 minutes.

cDNA Quantification Using qPCR

- 8 Thaw the GoTaq® Colorless Master Mix at room temperature. Vortex the Master Mix, then spin it briefly in a microcentrifuge to collect the material at the bottom of the tube.
- 9 Prepare one of the following reaction mixes on ice:

Component	Volume	Final Conc.
GOTaq® Colorless Master Mix, 2X	10 μ L	1X
Forward primer, 10 μ M (Jorge et al, 2017)	0.5 μ L	0.1 – 1.0 μ M

Reverse primer, 10 μ M (Jorge et al, 2017)	0,5 μ L	0,1 – 1,0 μ M
DNA template	5 μ L	< 250 ng
Nuclease-Free Water to	20 μ L	NA

- 10 Centrifuge the reactions in a microcentrifuge for 5 seconds.

- 11 Place the reactions in a thermal cycler that has been preheated to 95°C. Perform PCR using your standard parameters.

Step	Cycles	Temperature	Time
Denaturation	1	95°C	2 minutes
Denaturation	35	95°C	30 seconds
Annealing	55°C	30 seconds	
Extension	72°C	1 minute	
Final Extension	1	72°C	10 minutes