FEB 06, 2020

FIXATIVE FOR MOSQUITO LARVAE

PLOS One

Rochelly da Silva Mesquita¹, Wanderli Pedro Tadei¹

¹Laboratório de Malária e Dengue, Instituto Nacional de Pesquisas da Amazonia, Manaus, Brazil

Andrii Kyrylchuk

MATERIALS

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.9qrh5v6

External link:
<https://doi.org/10.1371/journal.pone.0227811>

Protocol Citation: Rochelly da Silva Mesquita, Wanderli Pedro Tadei 2020. FIXATIVE FOR MOSQUITO LARVAE. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.9qrh5v6>

MANUSCRIPT CITATION: Mesquita RdS, Kyrylchuk A, Grafova I, Kliukovskyi D, Bezudnyy A, Rozhenko A, Tadei WP, Leskelä M, Grafov A (2020) Synthesis, molecular docking studies, and larvicidal activity evaluation of new fluorinated neonicotinoids against *Anopheles darlingi* larvae. PLoS ONE 15(2): e0227811. doi: [10.1371/journal.pone.0227811](https://doi.org/10.1371/journal.pone.0227811)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Reagent	Quantity
Borax	5.0 g
Glycerol	2.5 mL
Formaldehyde (37... 40 % aq. soln.)	10.0 mL

Protocol status: Working

Created: Nov 24, 2019

Last Modified: Feb 06, 2020

PROTOCOL integer ID:
30193

- 1 Accurately weigh 5.0 g of borax and dissolve it in 100 mL of distilled water. Then, add 10 mL of the formaldehyde solution and 2.5 mL of glycerol, mix and bring to volume of 1000 mL with distilled water.