VERSION 2

DEC 31, 2019

Fluorescence Titering Assay for Lentivirus V.2

In 1 collection

Addgene The Nonprofit Plasmid Repository¹

¹Addgene

Addgene The Nonprofit Plasmid Repository
Addgene

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.bawtifen

External link:

<https://www.addgene.org/protocols/fluorescence-titering-assay/>

Protocol Citation: Addgene The Nonprofit Plasmid Repository 2019. Fluorescence Titering Assay for Lentivirus. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bawtifen>

MANUSCRIPT CITATION:

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Dec 31, 2019

Last Modified: Dec 31, 2019

PROTOCOL integer ID: 31411

Keywords: Fluorescence, Titering, Lentivirus, addgene

ABSTRACT

This protocol is for running a fluorescence titering assay for lentivirus. To see the full abstract and additional resources, visit the [Addgene protocol page](#).

Sample Data

Figure 1: Lenti-X 293T cells were transduced with a range of dilutions of 64108 pHAGE-TO-dCas9-3XmCherry. 72 hours post transduction, cells were assayed for mCherry expression using fluorescence microscopy.

GUIDELINES

Workflow Timeline

Day 0: Seed Lenti-X 293T cells (this cell line is optimized for production of lentiviral vectors)

Day 1: Transduce Cells

Day 2 (am): Remove media, replace with fresh media

Day 4-5: Count Fluorescent Cells

MATERIALS

Reagents

- DMEM high glucose
- L-alanyl-L-glutamine (or alternative stable glutamine)
- Heat-inactivated FBS
- Polybrene
- PBS pH 7.4 without calcium or magnesium (cations can affect the attachment of adherent cells)
- Microcentrifuge tubes
- 6-well dishes
- Pipettes
- Pipette tips
- Lentivirus Preparation

Equipment

- Biosafety cabinet
- Pipetman
- Pipettors
- Incubator
- Fluorescence microscope

Reagent Preparation

DMEM Complete: 10% v/v FBS and 4 mM L-alanyl-L-glutamine

- To a 500 mL bottle of DMEM high glucose, add 55 mL of heat inactivated FBS and 11 mL of 200 mM L-alanyl-L-glutamine. Store at 4 °C .

Note

Pro-Tip

Different brands and lots of FBS can promote or inhibit transfection. Test a variety of brands and lots of FBS to find one suitable with your protocols. FBS can be purchased already heat inactivated or it can be inactivated in the lab by heating to 56 °C for 00:30:00 .

SAFETY WARNINGS

See SDS (Safety Data Sheet) for safety warnings and hazards.

BEFORE START INSTRUCTIONS

Considerations Before You Start

- The health of the the target cell line is critical for obtaining accurate titers.
 - Check the cells for [mycoplasma](#) regularly
 - Do not over or under-grow your cells.
 - Thaw a new vial of cells after 20-30 passages.
- Do not add penicillin/streptomycin to the media.
- Titer will vary between cell lines.
- It is not recommended that lentiviral supernatants be subjected to multiple freeze-thaw cycles.

Seeding cells

- 1 Seed 75,000 cells into each well of a 6-well dish.
 - 1.1 Dilute 525,000 cells into 14 mL of DMEM complete.
 - 1.2 Mix well by pipetting or inverting.
 - 1.3 Aliquot 2 mL of cell suspension into each well of the 6-well dish.
- 2 Incubate the cells overnight.

Viral Titering

- 3 If using freshly collected virus, filter through a 0.45 µm polyethersulfone filter to remove cells and debris.

Note

Lentiviral titer can decrease during cycles of freeze-thaw. If you are freezing and aliquoting virus, it is recommended that you titer from the frozen stock to account for any loss in titer associated with freeze-thaw.

- 4 If using frozen virus, rapidly thaw the lentiviral aliquot at 37 °C by agitating in a warm water or bead bath.

- 5 Prepare dilutions of the lentivirus into DMEM complete containing 10 µg/mL polybrene.

Note

Note, this protocol was developed using low titer lentiviral vectors (10^5 TU/mL). If you anticipate that your viral stock will be higher titer consider additional dilutions.

Dilution	Volume of Lentivirus Stock (µL)	Volume of DMEM complete (µL)	Volume of 10mg/mL polybrene (µL)
1:10	150	1348.5	1.5
1:25	60	1438.5	1.5
1:50	30	1468.5	1.5
1:75	20	1478.5	1.5
1:100	15	1483.5	1.5

Mix the dilutions well

- 6 Gently aspirate media from the cells.
- 7 Add 1.5 mL of a viral dilution to each well (each well gets one dilution with one well left over).
- 8 Count the cells in the remaining well, a cell count is required for calculating the titer.
- 9 Incubate for 48:00:00 - 72:00:00 .
- 10 Gently aspirate media and replace with 1 mL of PBS.

Analysis

- 11 Calculate the percent of fluorescent-positive cells in each well.

Note

When calculating titer, only consider wells with less than 40% fluorescent-positive cells. Titering methods assume 1 integration event per cell. When the percentage exceeds 40% you risk counting cells with multiple integration events leading to underestimation of the true titer.

- 12 Calculate the transduction units per mL (TU/mL) using either the dilution factors (method 1) or the volume of virus (method 2):

12.1 Method 1: $\text{TU/mL} = (\text{Number of cells transduced} \times \text{Percent fluorescent} \times \text{Dilution Factor}) / (\text{Transduction Volume in mL})$

Note

Method 1 example: If the 1:100 well has 25% fluorescent cells and 150,000 cells were originally transduced, then there are
 $(150,000 \times 0.25 \times 100) / (1.5 \text{ mL}) = 2.5 \times 10^6 \text{ TU/mL}$

12.2 Method 2: $\text{TU/mL} = (\text{Number of cells transduced} \times \text{Percent fluorescent}) / (\text{Virus volume in mL})$

Note

Method 2 example: If 15 μL of virus added to 150,000 cells resulted in 25% fluorescent cells, then there are
 $(150,000 \times 0.25) / (0.015 \text{ mL}) = 2.5 \times 10^6 \text{ TU/mL}$

13 For a more accurate titer, take the average of multiple dilutions.