FEB 14, 2020

🌐 Fresh Frozen Mouse Brain Preparation (for Single Nuclei Sequencing)

Chuck Vanderburg¹, Carly Martin¹, Velina Kozareva¹,
Naeem Nadaf¹, Nehal Patel², Evan Macosko¹

¹Broad Institute of MIT and Harvard; ²Massachusetts General Hospital

The Single Cell Ninjas

Chuck Vanderburg

Broad Institute - Stanley Center

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.bcbrism6

External link:

<https://doi.org/10.1038/s41592-021-01264-7>

Protocol Citation: Chuck Vanderburg, Carly Martin, Velina Kozareva, Naeem Nadaf, Nehal Patel, Evan Macosko 2020. Fresh Frozen Mouse Brain Preparation (for Single Nuclei Sequencing).

protocols.io

<https://dx.doi.org/10.17504/protocols.io.bcbrism6>

MANUSCRIPT CITATION:

Biancalani T, Scalia G, Buffoni L, Avasthi R, Lu Z, Sanger A, Tokcan N, Vanderburg CR, Segerstolpe

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: Feb 07, 2020

Last Modified: Feb 14, 2020

PROTOCOL integer ID:
32849

- 1 Mice are anesthetized with 3% isoflourane prior to cardiac perfusion with ice cold HEPES-Sucrose Cutting Solution:

HEPES-Sucrose Cutting Solution				
	MW	final concentration (mM)	for 1L (in g or ml)	for 2L (in g or ml)
NaCl	58.44	110	6.43	12.86
HEPES	238.3	10	2.383	4.766
glucose	180.2	25	4.505	9.01
sucrose	342.3	75	25.6725	51.345
MgCl ₂ 6H ₂ O	203.31	7.5	7.5 ml (1M stock)	15 ml (1M stock)
KCl	74.56	2.5	2.5 ml (1M stock)	5 ml (1M stock)
adjusted to pH 7.4 with NaOH				

Store cutting solution at 4°C

- 2 Their head is removed by bisecting between c2 and c3 with large scissors. The brain is removed immediately as follows:
- 3 The cranial and facial skin and fat pads are peeled forward and the splenius capitis muscle is removed with a forceps exposure the dural sheath covering the fossa between the foramen magnum and the first cervical vertebra. A small pointed scissor is used to pierce the sheath in a rostral-caudal direction and then the 1 or 2 cervical vertebra are cut along the dorsal midline.
- 4 The same scissors are then inserted carefully 1-2mm into the foramen magnum with the lower blade pressing upward against the inner edge of the skull cavity. Proceed to snip the skull along the midline keeping constant gentle upward pressure so as not to cut or piece or otherwise damage the brain. Cut sagittally all the way up along the midline through the intra-orbital area over the olfactory bulbs.
- 5 Using a small toothed forceps, grab the midline edge of the bone covering the cerebellum and gently push the entire hemisphere of cut bone away from the midline. It will break, and only part of it will be removed from over the underlying brain tissue. Repeat the process until the bone from the foramen magnum up to the orbit is removed. Then do the other side. Be most careful of the cerebellar paraflocculus - that lobe nestles inside of a little fossa of the temporal bone along the lateral side of the cerebellum (ie, behind the ear) and we don't want to tear it off.

The bone around the cribiform plate and orbit is particularly thick - this must be carefully picked

- 6 away around the olfactory bulbs which are inside of it. Once the bulbs are fully exposed on the dorsal aspect they can be gently eased caudally using a blunt instrument like a small weighing spatula. As you release the olfactory bulbs, continue to “roll” the brain up and back from the inner rostral skull. The optic nerves will appear under the brain next - use the small scissors to cut them leaving the chiasma intact (this makes a great landmark later). Next comes the carotid artery - cut as well - then towards the posterior the basilar artery and some of the small cranial nerves appear. Try to cut vs. tear these if you can.
- 7 The entire operation thus far should have been less than ten minutes from start of anesthesia. If you are taking longer, you just need practice, don't waste your time using the RNA from practice brains.
- 8 Now the brain is ready to freeze. Wash in cold PBS to remove any hair, blood or misc. debris. Blot gently the excess PBS, not dry, but not swimming in liquid that will turn to ice in the next steps. Watch the movie.
- 9 Keep the brain frozen at -80 until use. We store whole mouse brains in a small 5 ml tube that has 2 ml of OCT frozen in the bottom (in advance). This OCT serves as a moisture reservoir that keeps the brain from sublimation and drying during long term storage. Desiccated brain tissue has poor RNA quality.