APR 02, 2020

Fruit fly potato culture media

Jorge Frias¹, Nelson Simões¹

¹CBA – Centre for Biotechnology of Azores, University of Azores, 9500-321 Ponta Delgada, Açores. Portugal.

Jorge Frias
Universidade dos Acores

ABSTRACT

Potato-based medium for culture of *Drosophila melanogaster*.

MATERIALS

- Distilled water;
- Dried yeast;
- White sugar;
- Instant mashed potato flakes;
- Agar;
- 95% ethanol;
- Ascorbic acid (Sigma - A92902);
- Methyl Paraben (Sigma – 47889);

BEFORE START INSTRUCTIONS

Preparation of 10 % Nipagin stock solution:

- 50 mL of 95% ethanol;
- 17.5 mL distilled water;
- 6.75 g of Methyl Paraben;

Note – This solution must be protected from light exposure.

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bei3jcgcn

Protocol Citation: Jorge Frias, Nelson Simões 2020. Fruit fly potato culture media. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bei3jcgcn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Apr 02, 2020

Last Modified: Apr 02, 2020

PROTOCOL integer ID:
35131

Keywords: culture media, fruit fly, *Drosophila*

1 Dissolve all following ingredients in 1.1 L of distilled water:

- 12.9 g Dried yeast;
- 1 g White sugar;
- 40 g Instant mashed potato flakes;
- 10 g Agar;
- 0.875 g Ascorbic acid;

- 2 Boil everything for 15 minutes.
- 3 Let it cool down to 60°C and add 8.5 mL of 10% Nipagin solution.
- 4 Mix well and fill about 1/4 of your preferred fruit fly vials or bottles with the medium.
- 5 On a clean bench hood, let the medium solidify in the vials with the cap open to avoid water condensation and store at 4°C for further usage.