

Apr 02, 2020

## Fruit fly potato culture media

DOI

[dx.doi.org/10.17504/protocols.io.bei3jcgn](https://dx.doi.org/10.17504/protocols.io.bei3jcgn)

Jorge Frias<sup>1</sup>, Nelson Simões<sup>1</sup>

<sup>1</sup>CBA – Centre for Biotechnology of Azores, University of Azores, 9500- 321 Ponta Delgada, Açores, Portugal.



Jorge Frias

Universidade dos Acores

---

OPEN  ACCESS



DOI: [dx.doi.org/10.17504/protocols.io.bei3jcgn](https://dx.doi.org/10.17504/protocols.io.bei3jcgn)

**Protocol Citation:** Jorge Frias, Nelson Simões 2020. Fruit fly potato culture media. [protocols.io](#)

<https://dx.doi.org/10.17504/protocols.io.bei3jcgn>

**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working

We use this protocol and it's working

**Created:** April 02, 2020

**Last Modified:** April 02, 2020

**Protocol Integer ID:** 35131

**Keywords:** culture media, fruit fly, Drosophila,

### Abstract

Potato-based medium for culture of *Drosophila melanogaster*.

## Materials

- Distilled water;
- Dried yeast;
- White sugar;
- Instant mashed potato flakes;
- Agar;
- 95% ethanol;
- Ascorbic acid (Sigma - A92902);
- Methyl Paraben (Sigma – 47889);

## Before start

Preparation of 10 % Nipagin stock solution:

- 50 mL of 95% ethanol;
- 17.5 mL distilled water;
- 6.75 g of Methyl Paraben;

Note – This solution must be protected from light exposure.

- 1 Dissolve all following ingredients in 1.1 L of distilled water:
  - 12.9 g Dried yeast;
  - 1 g White sugar;
  - 40 g Instant mashed potato flakes;
  - 10 g Agar;
  - 0.875 g Ascorbic acid;
- 2 Boil everything for 15 minutes.
- 3 Let it cool down to 60°C and add 8.5 mL of 10% Nipagin solution.
- 4 Mix well and fill about 1/4 of your preferred fruit fly vials or bottles with the medium.
- 5 On a clean bench hood, let the medium solidify in the vials with the cap open to avoid water condensation and store at 4°C for further usage.