AUG 09, 2018

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.sh4eb8w

Protocol Citation: Fatma Gomaa, Zhuhong Li, Roberto Docampo, Peter Girguis, Virginia Edgcomb 2018. G418 Kill curve protocol .
protocols.io
<https://dx.doi.org/10.17504/protocols.io.sh4eb8w>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Aug 09, 2018

Last Modified: Aug 09, 2018

PROTOCOL integer ID:
 14620

G418 Kill curve protocol

Fatma Gomaa¹, Zhuhong Li², Roberto Docampo², Peter Girguis¹, Virginia Edgcomb³

¹Harvard University; ²University of Georgia;

³Woods Hole Oceanographic Institution

Protist Research to Optimize Tools in Genetics (PROT-G)

Fatma Gomaa
 Harvard University

ABSTRACT

***Bodo saltans* kill curve protocol using G418 (Gentamicin)**

- Gentamycin exhibits toxicity toward both eukaryotic and prokaryotic cells by disrupting ribosome function, thereby blocking the elongation step in protein synthesis.
- G418 is most commonly used as a selection agent for eukaryotic cells genetically engineered to express a neomycin resistance gene (NeoR), which is encoded by either transposon Tn601 (903) or Tn5.
- The resistant cells grow in medium containing G418, and may be used to establish stably transfected cell lines as all the non-resistant cells die due to G418 toxicity, typically within 6 – 14 days.
- To establish kill curve assay for *saltans* various concentrations of G418 were tested. The concentration range from 1 to 50 µg/mL for selection of eukaryotic, and then higher concentrations can be used for maintaining stable transfected cell line.

Kill Curve Assay

1. Harvest the *B. saltans* cells from a culture that is at peak density ($1 - 3.0 \times 10^5$) by centrifugation at 800 x g for 5 min., discard the supernatant (medium).
1. Replace the growth medium with fresh medium containing 0 – 50 µg/mL. For each concentration, test in triplicate.
1. Using 6 wells plates, replace the medium in the wells every 3 – 4 days using fresh medium with the appropriate G418 concentration. Perform a daily visual inspection for evidence of toxicity, also cell count using hemocytometer.
1. Note that the optimal dose of G418 for selection is the lowest one for which all

cells have died after one week. A low dose is the concentration which has minimal effects on cells after 2 weeks of antibiotic selection. Conversely, a high dose is a concentration which is highly toxic to cells within 2-3 days of starting antibiotic selection.

5- The results indicate that a G418 concentration of 2 µg/mL kill the entire *B. saltans* population in 12 days which is considered enough time for selecting the resistant cells after transfection.

G418 Sensitivity for *B. saltans*

G 4 1 8 (µ g / m l)	C e l l s u r v i v a l (d a y s)
5 0	2
2 0	2
1 5	3
1 0	4
5	6
3	8
2	1 2
1	> 1 6
0	> 1 6

G 4 1 8 (μ g / m l)	C e l l s u r v i v a l (d a y s)
---	--

Cell Selection

1. In all of our plasmids constructs we included the *NeoR* gene (Neomycine resistant gene) to select our transfected cells.
2. Twenty-four hours after electroporation, the growth medium of transfected cells were supplemented with G418 antibiotic Solution (2 μg/mL) based on the kill curve results (above).
3. Replace the G418-containing medium every 3 – 5 days and examine cells for visual toxicity. Most non-transfected (non-resistant) cells will die within 10- 12 days, leaving the transfected cells to expand.
2. Once cells grow to high confluence, they may be maintained, or frozen as a polyclonal line or plated by limited dilution to select for single clones.

ATTACHMENTS

Kill
curve_B_saltans_FG.docx