VERSION 4
MAR 02, 2021

General bacteria and archaea 16S-rRNA (515Fmod-806Rmod) for Illumina amplicon sequencing V.4

Roey Angel¹, Eva Petrova¹

¹Soil and Water Research Infrastructure

SoWa RI Anaerobic and Molecular Microbiology (public)

Tech. support email: eva.petrova@bc.cas.cz

Roey Angel

Soil and Water Research Infrastructure

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.bsxanfie

Protocol Citation: Roey Angel, Eva Petrova 2021. General bacteria and archaea 16S-rRNA (515Fmod-806Rmod) for Illumina amplicon sequencing.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.bsxanfie> Version created by [Roey Angel](#)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Mar 02, 2021

Last Modified: Mar 02, 2021

PROTOCOL integer ID:
47810

Keywords: PCR, 16S rRNA, SSU rRNA, Amplicon sequencing, Illumina sequencing, Barcoded sequencing, Targeted metagenomics, Microbiome

ABSTRACT

Universal 16S rRNA probe-based-qPCR assay for bacteria.

The primers target the V4 region of the 16S rRNA gene and were specifically designed for Illumina amplicon sequencing. The original primers were designed by Caporaso *et al.* (2012) and modified by Walters *et al.* (2015). For barcoding, we use the [Fludigm Access Array](#) for barcoding the sample and therefore the primers are synthesized with the CS1 and CS2 regions.

CITATION

Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, Owens SM, Betley J, Fraser L, Bauer M, Gormley N, Gilbert JA, Smith G, Knight R (2012). Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms.. The ISME journal.

LINK

<https://doi.org/10.1038/ismej.2012.8>

CITATION

Walters W, Hyde ER, Berg-Lyons D, Ackermann G, Humphrey G, Parada A, Gilbert JA, Jansson JK, Caporaso JG, Fuhrman JA, Apprill A, Knight R (2015). Improved Bacterial 16S rRNA Gene (V4 and V4-5) and Fungal Internal Transcribed Spacer Marker Gene Primers for Microbial Community Surveys.. mSystems.

LINK

[10.1128/msystems.00009-15](https://doi.org/10.1128/msystems.00009-15)

MATERIALS

STEP MATERIALS

- ✕ Agarose Merck MilliporeSigma (Sigma-Aldrich) Catalog #A9539
- ✕ GeneRuler DNA Ladder Mix Thermo Fisher Scientific Catalog #SM0331
- ✕ DNA Gel Loading Dye (6X) Thermo Fisher Scientific Catalog #R0611
- ✕ TAE buffer (50x), molecular biology grade Serva, Germany Catalog #4254901
- ✕ Primer: 515Fmod_CS1 Elisabeth Pharmacon
- ✕ Primer: 806mod_CS2 Elisabeth Pharmacon
- ✕ DreamTaq Green DNA Polymerase (5 U/μL) Thermo Fisher Scientific Catalog #EP0712
- ✕ dNTP Set (100 mM each) Contributed by users Catalog #BR0600601
- ✕ PCR H2O Top Bio Catalog #P040
- ✕ Bovine Serum Albumin (BSA) Thermo Fisher Scientific Catalog #B14

Primers

1

Name	Direction	Sequence ¹	Target region ²
515Fmod_CS1	Forward	ACA CTG ACG ACA TGG TTC TAC AGT GYC AGC MGC CGC CGT AA	515-533
806Rmod_CS2	Reverse	TAC GGT AGC AGA GAC TTG GTC TGG ACT ACN VGG GTW TCT AAT	787-806

1. CS + primer sequence (in bold)
2. Relative to *E. coli* SSU rRNA gene

✕ Primer: 515Fmod_CS1 Elisabeth Pharmacon

✕ Primer: 806mod_CS2 Elisabeth Pharmacon

PCR reaction

2 Prepare the following master mixture On ice .

Do not forget to prepare some additional mixture for the negative (NTC) and positive controls, and to account for pipetting errors.

A	B	C	D
Reagent	Final. conc.	1 tube (25 μ l)	100 reactions (96-well plate; μ l)
PCR H ₂ O		17.525	1752.5
10X DreamTaq Green Buffer	1X	2.5	250
dNTP (2 mM each)	0.2 mM	2.5	250
BSA (20 μ g μ l ⁻¹)	80 ng μ l ⁻¹	0.1	10
515Fmod-CS1	0.2 μ M	0.625	62.5
806Rmod-CS2	0.2 μ M	0.625	62.5
DreamTaq Green DNA Polymerase	0.625 U	0.125	12.5
Final volume		24	2400

DreamTaq Green DNA Polymerase (5 U/ μ L) Thermo Fisher Scientific Catalog
#EP0712

dNTP Set (100 mM each) Contributed by users Catalog
#BR0600601

PCR H₂O Top Bio Catalog
#P040

Bovine Serum Albumin (BSA) Thermo Fisher Scientific Catalog
#B14

3 Vortex and spin down 00:00:03

3s

4 Distribute 24 μ L of the mixture to each tube and add 1 μ L of template DNA or cDNA

PCR reaction

3s

5 Run the following PCR program:

17m 15s

1. 94 °C 00:05:00
2. x 28 {
 - 2.1 94 °C 00:00:45
 - 2.2 52 °C 00:00:45
 - 2.3 72 °C 00:00:45
3. 72 °C 00:10:00
4. 4 °C hold

Evaluate PCR products on an agarose gel

40m

6 Prepare a 1.5% agarose gel by mixing:

100 mL
TAE

1.5 g
agarose

Heat in the microwave until dissolved and pour into a gel frame.

Place solid gel into an electrophoresis bath filled with TAE buffer.

Agarose Merck MilliporeSigma (Sigma-Aldrich) Catalog
#A9539

GeneRuler DNA Ladder Mix Thermo Fisher Scientific Catalog
#SM0331

DNA Gel Loading Dye (6X) Thermo Fisher Scientific Catalog
#R0611

TAE buffer (50x), molecular biology grade Serva, Germany Catalog
#4254901

7 Mix up to 5 µL of the PCR reaction sample with 1 µL of loading dye and load the sample into a well.
In addition load 5 µL of DNA ladder mix (80-10,000 bp) into an empty well, as a marker.

8 Run the gel at 110V, 265mA for approx. 00:40:00

40m

-
- 9** Stain gel for at least 40min in an Ethidium bromide TAE bath (or any other DNA stain).
 - 10** Visualise the gel using a gel documentation system.