NOV 14, 2020

Generation of Combinatorial CRISPR Libraries

Nature Communications

David Adams¹, Nicky Thompson. nt4@sanger.ac.uk. ¹

¹Wellcome Sanger Institute

David Adams

DISCLAIMER

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.bpqhmmmt6

External link:

<https://doi.org/10.1038/s41467-021-21478-9>

Protocol Citation: David Adams, Nicky Thompson. nt4@sanger.ac.uk. 2020. Generation of Combinatorial CRISPR Libraries. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bpqhmmmt6>

MANUSCRIPT CITATION:

Thompson NA, Ranzani M, Weyden Lvd, Iyer V, Offord V, Droop A, Behan F, Gonçalves E, Speak A, Iorio F, Hewinson J, Harle V, Robertson H, Anderson E, Fu B, Yang F, Zagnoli-Vieira G, Chapman P, Velasco-Herrera MDC, Garnett MJ, Jackson SP, Adams DJ, Combinatorial CRISPR screen identifies fitness effects of gene paralogues. Nature Communications doi: [10.1038/s41467-021-21478-9](https://doi.org/10.1038/s41467-021-21478-9)

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to protocols.io is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with protocols.io, can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Nov 14, 2020

Last Modified: Nov 14, 2020

PROTOCOL integer ID:
44521

Summary

- 1 This protocol is a step by step protocol based on the method originally described by: Vidigal, J.A. & Ventura, A. Rapid and efficient one-step generation of paired gRNA CRISPR-Cas9 libraries. *Nature Communications* **6**, 8083 (2015).

Throughout:

All PCR clean-ups done with Monarch PCR & DNA cleanup kit

<https://www.neb.com/products/t1030-monarch-pcr-dna-cleanup-kit-5-ug>

All gels run with Sybr green, not ethidium bromide, and no gels exposed to UV light to avoid DNA damage.

Vector Preparation

2 Backbone digestion and preparation (lenti-guide puro)

Digest 5-10ug of your vector of choice:

1. lenti-guide puro 10ug (Addgene #52963)
2. 10X buffer 3.1 10ul
3. BsmBI 10000 U/ml 8ul
4. H2O to 100ul

Incubate @ 55°C overnight

Dephosphorylation

<https://www.neb.com/protocols/1/01/01/vector-dephosphorylation-protocol>

- 49ul of digestion
- Antarctic phosphatase 5ul
- Antarctic phosphatase buffer 10X 6ul

37C for 45'

Then 5' 80C

Then 4C

Gel purification: take the 8.3kb band.

PCR and oligos

3 **PCR amplification of oligonucleotides**

Supplied by TWIST bioscience

Diluted library to final concentration of 0.0045ng/ul

Do 14xPCR – each PCR 50ul

<https://www.neb.com/protocols/1/01/01/pcr-protocol-m0530>

	Rege nt amou nt for 14x reacti ons
Oligo	14
HF phusion buffer	140
dNTP	14
F primer	17.5
R primer	17.5
Phusion HF polymerase	7
H2O	490
	700

PCR program

- a. 98C 30'
- i. 98 10"
- ii. 68 C° 35'
- iii. 72 C 30" 16X
- b. 72C 10'
- c. 4C 4ever

PCR cleanup and elute in 30ul

Digestion of pDonor_SU6

Digest 5ug of pDonor_sU6 (plasmid #741) with *BbsI*

Plasmid 5ug
Buffer 2.1 10X 5ul
Enzyme BbsI 10000 U/ml 3ul
H2O up to 50ul total

For 37C for three hours
Gel purification: take the 415bp band

Gibson Ligation

<https://www.neb.com/products/t1030-monarch-pcr-dna-cleanup-kit-5-ug>

Donor fragment 405ng
Oligo amplicon 432ng
2xGibson MM 30 ul (NEBuilder® HiFi DNA Assembly Master Mix, E2621S)
H2O to 60 ul

Incubate @ 50°C for 2 hours

Nuclease Digestion

<http://www.epibio.com/enzymes/nucleases-glycosylases-dna-binding-proteins/dna-exonucleases/plasmid-safe-atp-dependent-dnase?details>

Add to each Gibson reaction

- 10x Plasmid Safe Buffer 9 ul
- ATP (25mM) 9 ul
- Plasmid Safe nuclease 3 ul
- H2O 9 ul

incubate @ 37°C for 1h

Clean up using PCR purification kit and elute in 50ul water

Digestion of Gibson product

Digest with *Bbsl*
DNA 50ul
Buffer 2.1 6ul
Bbsl 4ul

Incubate @ 37°C for 2.5h
Run digestion on 2.5% gel and cut ~480 bp band
Gel extract and elute the completed insert in 30ul

Ligation

<https://www.neb.com/products/m2200-quick-ligation-kit>

Neb Quick ligation kit; M2200S; scale up as appropriate

	Amount	
Digested&purified lenti-guide puro	25ng	
sgRNA insert Bbsl purified	X	
ligase buffer	5ul	
Ligase	0.5ul	
H2O to 10.5ul		

Incubated at 25C for 15minutes (thermocycler)
Prior to library creation controls done with vector only (lenti-guide puro) to assess background.
For library used final volume of 300ul for ligation
PCR purification and elution in 30ul water (for library)

Transformation

4 Transformation

<https://www.neb.com/products/c3020-neb-10-beta-electrocompetent-e-coli>

25ul bacteria/2.5ul ligation product/transformation; electroporated as per website protocol. 10 electroporations. 1 hour in pre-warmed SOC shaking.

Plated at dilutions (1:1000; 1:10000) onto warmed ampicillin plates. The remainder inoculated into ampicillin containing LB broth 3 litres at 100ug/ml and shaken at 225rpm/37C overnight.

Following morning pooled spun down in Avanti centrifuge (6000g for 10 minutes)

Bacterial pellet processed using maxi/mega prep (Qiagen) ~1.5g bacteria/mega prep