VERSION 1
OCT 17, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.8d3hs8n

Protocol Citation: Vinca Seiler, René Inckemann 2019. Golden Gate lvl 0.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.8d3hs8n>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Oct 17, 2019

Last Modified: Oct 17, 2019

PROTOCOL integer ID:
28827

🌐 Golden Gate lvl 0 V.1

Vinca Seiler¹, René Inckemann¹

¹iGEM Team Marburg 2019

Igem Marburg
iGEM Marburg 2019

ABSTRACT

Golden Gate reaction protocol for lvl 0

MATERIALS

MATERIALS

✂ BsmBI - 1,000 units **New England Biolabs Catalog #R0580L**

✂ T4 DNA Ligase **New England Biolabs Catalog #M0202**

✂ Esp3I **New England Biolabs Catalog # R0734L**

✂ 10X NEB T4 DNA ligase buffer **New England Biolabs**

1 0.5 µL of DNA insert (60 ng/µl)

2 0.5 µL of entry Vector (15 ng/ µL)

3 1 µL T4 DNA Ligase buffer (NEB)

4 0.5 µL T4 DNA Ligase (NEB)

5 0.5 µL EspI3 (NEB)

6 Water to 10 µL

Thermocycler conditions

7

37 °C 00:20:00

8

37 °C 00:01:30

9

16 °C 00:03:00

10 Cycle step 8 and 9 5-10x

11 50 °C 00:05:00

12 80 °C 00:10:00

Transformation

13 Add 2 µL - 5 µL of each assembly reaction to 50 µL competent cells.

14 Cells should be recovered for 01:00:00 (Amp) to 02:00:00 (Kan, Chloramphenicol).