Growth curve for *Chlamydomonas reinhardtii*

Joao Vitor Molino¹

¹Ronin Institute

Joao Vitor Molino
Ronin Institute

DISCLAIMER

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](https://dx.doi.org/10.17504/protocols.io.bpvbm2n) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](https://dx.doi.org/10.17504/protocols.io.bpvbm2n), can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

ABSTRACT

This protocols describe the steps required for obtain a growth curve of *Chlamydomonas reinhardtii* and fluorescent protein expression of mVenus and mCherry.

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bpvbm2n

Protocol Citation: Joao Vitor Molino 2020. Growth curve for *Chlamydomonas reinhardtii*. [protocols.io](https://dx.doi.org/10.17504/protocols.io.bpvbm2n) <https://dx.doi.org/10.17504/protocols.io.bpvbm2n>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Nov 19, 2020

Last Modified: Nov 19, 2020

PROTOCOL integer ID:
44675

Keywords: Growth curve, *Chlamydomonas*, Absorbance, 96 well plate

Material

- 1
 - TAP media or other (How to prepare TAP media [here](#))
 - Erlenmeyer flask
 - Orbital shaker
 - Light source
 - 96 well plate, Black Frame, Clear bottom (Ex:

⊗ IsoPlate™ - 96F (Black Frame & Clear well 96-well) **Perkin Elmer Catalog #6005020**

)

Plate reader Settings

- 2 Reading are performed in a Black 96 well plate with clear bottom.

Absorbance set to 750 nm .

Fluorescence set as the Table below.

	Excitation (nm)	Emission (nm)	Gain	Optics position
Chlorophyll	440	680	70	bottom
mVenus	500	530	120	bottom
mCherry	583	613	150	bottom

Inoculum

2w 5d

- 3 Inoculate 1 mL of the cells from a culture in stationary phase (120:00:00 to 168:00:00) in an 100 mL erlenmeyer flask containing 50 mL of TAP media.

2w 5d

1. Place the flask in a 150 rpm, 25°C, 1cm of orbit with 80 $\mu\text{mol}/\text{m}^2\text{s}$ of incident white light (60 $\mu\text{mol}/\text{m}^2\text{s}$ to 120 $\mu\text{mol}/\text{m}^2\text{s}$ works)
2. Take a initial sample 100 μL of culture and measure it in the plate reader according to the settings above.
3. During 168:00:00 take samples at least once a day.
4. The final culture can be used for further test, as dry cell weight (DCW) determination.

Note

Frequent sampling increase data quality, but it is advice to not remove more than 10% of culture in sampling during the entire procedure. Technical replicates are advice for each time point.

All culturing conditions are set initially, and can be change accordingly to the experiment goal.

Example of DCW protocol.

Protocol

NAME

Dry cell weight by centrifugation

CREATED BY

Joao Vitor Molino

PREVIEW

- 3.1**
- Analytical balance with high precision (*The higher the precision the better. For example a balance with a 0.1mg readability, could account to approximately 10% error alone in a measurement of 1mL sample of a culture at 1g/L*)
 - Microcentrifugal tubes
 - Microcentrifuge
- 3.2**
1. Label microcentrifugal tubes
 2. Dry the tubes at 90 °C , Overnight
 3. Cool tubes at Room temperature for 00:30:00
 4. Record the weight of the tubes
- 3.3**
1. Harvest 2 mL of culture in a previously weighted tube
 2. Centrifuge the sample at 20000 rcf, 25°C, 00:01:00
 3. Carefully remove the supernatant by pipetting
 4. Wash the cells with ddH2O, and centrifuge the sample at 20000 rcf, 25°C, 00:01:00
 5. Carefully remove the supernatant by pipetting

-
6. Dry the tubes at 90 °C , Overnight
 7. Cool tubes at Room temperature for 00:30:00
 8. Record the weight of the tubes
 9. Subtract the initial tube weight to achieve the dry cell weight