


Nov 19, 2020

Growth curve for *Chlamydomonas reinhardtii*

DOI

dx.doi.org/10.17504/protocols.io.bpvbmn2n



Joao Vitor Molino¹

¹Ronin Institute



Joao Vitor Molino

Ronin Institute, University of California, San Diego

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.bpvbmn2n

Protocol Citation: Joao Vitor Molino 2020. Growth curve for *Chlamydomonas reinhardtii*. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bpvbmn2n>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: November 19, 2020

Last Modified: November 19, 2020

Protocol Integer ID: 44675

Keywords: Growth curve, *Chlamydomonas*, Absorbance, 96 well plate,



Disclaimer


DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to **protocols.io** is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with **protocols.io**, can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

Abstract

This protocols describe the steps required for obtain a growth curve of *Chlamydomonas reinhardtii* and fluorescent protein expression of mVenus and mCherry.

Protocol materials

 IsoPlate™ - 96F (Black Frame & Clear well 96-well) **Perkin Elmer Catalog #6005020**

Material

- 1
 - TAP media or other (How to prepare TAP media [here](#))
 - Erlenmeyer flask
 - Orbital shaker
 - Light source
 - 96 well plate, Black Frame, Clear bottom (Ex:



IsoPlate™ - 96F (Black Frame & Clear well 96-well) Perkin Elmer Catalog #6005020

)

Plate reader Settings

- 2 Reading are performed in a Black 96 well plate with clear bottom.

Absorbance set to 750 nm .

Fluorescence set as the Table below.

	Excitation (nm)	Emission (nm)	Gain	Optic s positi on
Chlorophy II	440	680	70	botto m
mVenus	500	530	120	botto m
mCherry	583	613	150	botto m







Inoculum

2w 5d

- 3 Inoculate 1 mL of the cells from a culture in stationary phase (120:00:00 to 168:00:00) in an 100 mL erlenmeyer flask containing 50 mL of TAP media.

2w 5d



1. Place the flask in a  150 rpm, 25°C ,1cm of orbit with  80 $\mu\text{mol}/\text{m}^2\text{s}$ of incident white light ( 60 $\mu\text{mol}/\text{m}^2\text{s}$ to  120 $\mu\text{mol}/\text{m}^2\text{s}$ works)
2. Take a initial sample  100 μL of culture and measure it in the plate reader according to the settings above.
3. During  168:00:00 take samples at least once a day.
4. The final culture can be used for further test, as dry cell weight (DCW) determination.

Note

Frequent sampling increase data quality, but it is advice to not remove more than 10% of culture in sampling during the entire procedure. Technical replicates are advice for each time point.
All culturing conditions are set initially, and can be change accordingly to the experiment goal.

Example of DCW protocol.

Protocol



NAME

Dry cell weight by centrifugation

CREATED BY

Joao Vitor Molino

PREVIEW

- 3.1
 - Analytical balance with high precision (*The higher the precision the better. For example a balance with a 0.1mg readability, could account to approximately 10% error alone in a measurement of 1mL sample of a culture at 1g/L*)
 - Microcentrifugal tubes
 - Microcentrifuge
- 3.2
 1. Label microcentrifugal tubes



2. Dry the tubes at 90 °C , Overnight

3. Cool tubes at Room temperature for 00:30:00

4. Record the weight of the tubes

3.3

1. Harvest 2 mL of culture in a previously weighted tube

2. Centrifuge the sample at 20000 rcf, 25°C, 00:01:00

3. Carefully remove the supernatant by pipetting

4. Wash the cells with ddH₂O, and centrifuge the sample at

20000 rcf, 25°C, 00:01:00

5. Carefully remove the supernatant by pipetting

6. Dry the tubes at 90 °C , Overnight

7. Cool tubes at Room temperature for 00:30:00

8. Record the weight of the tubes

9. Subtract the initial tube weight to achieve the dry cell weight