OCT 06, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.7zghp3w

External link:
<https://doi.org/10.1371/journal.pone.0225863>

Protocol Citation: Norfitriah Mohamed Sohaimi, Mohd Hair Bejo, Abdul Rahman Omar, Aini Ideris, Nurulfiza Mat Isa 2019. Haematoxylin and eosin staining. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.7zghp3w>

MANUSCRIPT CITATION:
 Sohaimi NM, Bejo MH, Omar AR, Ideris A, Isa NM (2019) Molecular characterization of fowl adenovirus isolate of Malaysia attenuated in chicken embryo liver cells and its pathogenicity and immunogenicity in chickens. PLoS ONE 14(12): e0225863. doi: [10.1371/journal.pone.0225863](https://doi.org/10.1371/journal.pone.0225863)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Haematoxylin and eosin staining

PLOS One

Norfitriah Mohamed Sohaimi¹, Mohd Hair Bejo¹, Abdul Rahman Omar¹, Aini Ideris¹, Nurulfiza Mat Isa²

¹Faculty of Veterinary Medicine, Universiti Putra Malaysia, Serdang, Selangor, Malaysia.;

²Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, Serdang, Selangor, Malaysia.

Norfitriah Mohamed Sohaimi

MATERIALS

MATERIALS

Dulbecco's Modified Eagle's Medium (DMEM) Sigma Aldrich Catalog #D5796

For detection of intranuclear inclusion bodies, primary CEL cells were grown on sterile coverslips [Adair, 1978]. Briefly, the growth medium was aspirated out from confluent monolayer and washed twice with sterile medium free serum prior to inoculation of 0.1mL viral supernatant. Culture was added with maintenance medium, DMEM and 2% FBS and kept for 48 hours incubation, harvested and fixed in 10% buffered formalin for 5 minutes and stained with haematoxylin and eosin (HE).

Protocol status: Working
We use this protocol and it's
working

Created: Oct 06, 2019

Last Modified: Oct 06, 2019

PROTOCOL integer ID:
28424