



Sep 16, 2022

Hamstring muscle architecture assessed sonographically using wide field of view: a reliability study

DOI

<https://dx.doi.org/10.17504/protocols.io.5qpvorxjdv4o/v1>

Kevin Cronin¹

¹University College Dublin



Kevin Cronin

University College Dublin

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.5qpvorxjdv4o/v1>

Protocol Citation: Kevin Cronin 2022. Hamstring muscle architecture assessed sonographically using wide field of view: a reliability study. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.5qpvorxjdv4o/v1>

License: This is an open access protocol distributed under the terms of the **[Creative Commons Attribution License](#)**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: September 07, 2022

Last Modified: September 16, 2022

Protocol Integer ID: 69688

Keywords: reliability study hamstring injury, hamstring muscle architecture, retest minimum detectable difference of the hamstring muscle, architecture of the hamstring muscle, vivo hamstring muscle architecture, hamstring muscle, hamstring strain injury, risk factor for hamstring strain injury, muscle architecture, ultrasounds field of view, view ultrasound, ultrasounds field, previous ultrasound technique, ultrasound, various medical imaging technique, magnetic resonance imaging, wide field of view

Abstract

Hamstring injuries are very common in field sports. Muscle architecture has been suggested as a risk factor for hamstring strain injury. Various medical imaging techniques (Magnetic Resonance Imaging and Ultrasound) have been developed to assess muscle architecture. Ultrasound is often used to assess in vivo hamstring muscle architecture. The architecture of the hamstring muscles often extends outside the ultrasounds field of view. Previous ultrasound techniques used often generate questionable results. This study aims to describe the reliability of large field of view ultrasound and test – retest minimum detectable difference of the hamstring muscles

Attachments



[Protocol.docx](#)

1.3MB

