VERSION 1

FEB 26, 2021

DOI:
dx.doi.org/10.17504/protocols.io.bss7nehn

External link:
<https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0253830>

Collection Citation: Ashley Jones, Cynthia Torkel, David Stanley, Jamila Nasim, Justin Borevitz, Benjamin Schwessinger 2021. High-molecular weight DNA extraction, clean-up and size selection for long-read sequencing. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bss7nehn>

MANUSCRIPT CITATION:
 Jones A, Torkel C, Stanley D, Nasim J, Borevitz J, Schwessinger B (2021) High-molecular weight DNA extraction, clean-up and size selection for long-read sequencing. PLoS ONE 16(7): e0253830.
doi.org/10.1371/journal.pone.0253830

License: This is an open access collection distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

High-molecular weight DNA extraction, clean-up and size selection for long-read sequencing V.1

PLOS One ✓ Peer-reviewed method

Ashley Jones¹, Cynthia Torkel¹, David Stanley¹, Jamila Nasim¹, Justin Borevitz¹, Benjamin Schwessinger¹

¹Research School of Biology, Australian National University, Canberra, ACT, Australia

High molecular weight DNA extraction from all kingdoms

PLOS ONE Lab Protocols

Ashley Jones
 Australian National University

ABSTRACT

Rapid advancements in long-read sequencing technologies have transformed sequencing read lengths from bps to Mbps, which has enabled chromosome-scale genome assemblies. However, read lengths are now becoming limited by the extraction of pure high-molecular weight DNA suitable for long-read sequencing, which is particularly challenging in plants and fungi. To overcome this, we present a protocol collection; high-molecular weight DNA extraction, clean-up and size selection for long-read sequencing. We optimised a gentle magnetic bead based high-molecular weight DNA extraction, which is presented here in detail. The protocol circumvents spin columns and high-centrifugation, to limit DNA fragmentation. The protocol is scalable based on tissue input, which can be used on many species of plants, fungi, reptiles, insects and bacteria. It is also cost effective compared to kit-based protocols and hence applicable at scale at low resource settings. An optional sorbitol wash is listed and is highly recommended for plant and fungal tissues. To further remove any remaining contaminants such as phenols and polysaccharides, optional DNA clean-up and size selection strategies are given. This protocol collection is suitable for all common long-read sequencing platforms, such as technologies offered by PacBio and Nanopore. Using these protocols, sequencing on the Oxford Nanopore MinION can achieve read length N50 values of 30-50 kb, with reads exceeding 200 kb and outputs ranging from 15-30 Gbp. This has been routinely achieved with eucalypts, acacias, rice, the meda, wheat, wheat rusts, various other fungi, geckos, skinks, ticks, ladybird beetles, caterpillars and E. coli.

Protocol status: Working
We use this protocol routinely across multiple genera.

Created: Feb 26, 2021

Last Modified: Feb 26, 2021

COLLECTION integer ID:
47679

FILES

Protocol

NAME
Scalable high-molecular weight DNA extraction for long-read sequencing

VERSION 1

CREATED BY

Ashley JonesAustralian National University

OPEN →

Protocol

NAME
Sorbitol washing complex homogenate for improved DNA extractions

VERSION 1

CREATED BY

Ashley JonesAustralian National University

OPEN →

Protocol

NAME
DNA clean-up and size selection for long-read sequencing

VERSION 3

CREATED BY

Ashley JonesAustralian National University

OPEN →