OCT 24, 2020

His-tag purification

Andreea S¹

¹University of Groningen

iGEM Groningen 2020

a.stan.6

ABSTRACT

His tag purification uses the technique of immobilised metal affinity chromatography. In this technique, transition metal ions are immobilized on a resin matrix using a chelating agent such as iminodiacetic acid. It has been studied that among amino acids constituting proteins, histidine is strongly involved in the coordinate bond with metal ions. Therefore, if a number of histidines are added to the end of the protein by genetic engineering, the affinity of the protein for the metal ion is remarkably increased and the basic idea is that purification can be easily carried out. When a protein having a His tag is brought into contact with a carrier on which a metal ion such as nickel is immobilized, the histidine residue chelates the metal ion and binds to the carrier. Since other proteins do not bind to the carrier, they can be washed off with a buffer. Thereafter, it is possible to recover the protein having the His tag with high purity.

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bnw5mfg6

Protocol Citation: Andreea S 2020. His-tag purification . **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bnw5mfg6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Other
 The protocol is developed based on literature and has not been tested yet.

Created: Oct 24, 2020

Last Modified: Oct 24, 2020

PROTOCOL integer ID:
 43709

Separate proteins from soil matrix

1 Collect soil samples of 5 g

- 2 Extract total proteins using NoviPure Soil Protein Kit or other commercially available kits for total protein soil extraction

His-tag separation of NLP

1h

- 3 Wash the Ni^{2+} -sepharose column material with 12 CVs of MQ and 4 CVs of Column Wash Buffer (10 millimolar (mM) imidazole, KPi 50 millimolar (mM) pH 7, NaCl 200 millimolar (mM)).

Note

Use ± 0.5 ml of Ni^{2+} -sepharose column material per 10 mg of total protein.

- 4 Apply the sample, add imidazole (10mM final concentration) and the washed Ni^{2+} -sepharose column material. Nutate in at 4 °C for 01:00:00

1h

- 5 Pour column, collect flow through to apply on SDS gel.

- 6 Wash column with 20 CVs of Wash Buffer (50 millimolar (mM) imidazole, KPi 50 millimolar (mM) pH 7, NaCl 200 millimolar (mM)).

- 7 Elute protein with Elution Buffer (500 millimolar (mM) imidazole, KPi 50 millimolar (mM) pH 7, NaCl 200 millimolar (mM)) in 200 μL fractions. Check elution fractions Absorbance by NanoDrop.

Run an SDS gel to check purification:

8

- Soil suspension & Flow through: dilute 15x, apply 5 μL
- Wash: dilute 1.25x, apply 10 μL
- Elution fractions: dilute to ± 0.2 mg/ml, apply 5 μL