FEB 13, 2019

OPEN ACCESS

DOI:

[dx.doi.org/10.17504/protocols.io.vg6e3ze](https://doi.org/10.17504/protocols.io.vg6e3ze)

External link:

<https://doi.org/10.1371/journal.pone.0210123>

Protocol Citation: Paloma Varela 2019. Holo-ZitRMG structure determination by x-ray crystallography. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.vg6e3ze>

MANUSCRIPT CITATION:

Varela PF, Velours C, Aumont-Niçaise M, Pineau B, Legrand P, Poquet I (2019) Biophysical and structural characterization of a zinc-responsive repressor of the MarR superfamily. *PLoS ONE* 14(2): e0210123. doi: [10.1371/journal.pone.0210123](https://doi.org/10.1371/journal.pone.0210123)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Holo-ZitRMG structure determination by x-ray crystallography

PLOS One

Paloma
Varela¹

¹Institute for Integrative Biology of the Cell, CEA, CNRS, Université Paris-Saclay, Gif-sur-Yvette, France

Paloma Varela

MATERIALS

Holo-ZitRMG protein was subjected to crystallization assays using sparse matrix screens in a robotic system (Nano-Robot Cartesian) (crystallography platform, LEBS/IMAGIF). In preliminary experiments crystals were first obtained using ZitRMG tetrameric form by vapor diffusion in 0.2 M ammonium acetate, 0.1 M sodium citrate tribasic dihydrate (pH 5.6) and 30 % polyethyleneglycol 4,000. They diffracted to a resolution of less than 6 Å. These crystals belong to $P4_1$ space group with cell parameters of $a = b = 129$ Å; $c = 88$ Å; $a = b = c = 90^\circ$. Crystallization conditions were then optimized using the dimeric form. New X-ray data were collected at the zinc peak absorption edge up to 3.7 Å resolution. Because molecular replacement with different MarR protein coordinates did not work, after data processing using XDS. we used Zn anomalous signal to solve the structure by SAD method using PHASER. Eight molecules were found in the asymmetric unit, with a solvent content of 60 %, and two zinc atoms were found per protomer. Phases were improved and extended by NCS averaging and solvent flattening using PARROT. Refinement was done using BUSTER-TNT. To help building this low-resolution model, *B. subtilis* OhrR structure was used. From a poly-Ala chain, we could identify the extended wHTH DNA-binding domain, but most of the lateral chains were not visible. New data collection recorded on a Pilatus 6M detector on the PROXIMA1 beamline at SOLEIL synchrotron went up to 2.8 Å resolution. An anisotropic correction factor was applied STARANISO server (<http://staraniso.globalphasing.org/>) to sharpen the data in the weaker-diffracting direction. Structure was determined by molecular replacement using AdcR (3TGN) as model. The high values of average B-factors of the final structure can be explained by the crystal anisotropy. All structure figures are created using The PyMOL Molecular Graphics System, Version 1.8 Schrödinger, LLC.

Created: Nov 09, 2018

Last Modified: Feb 13, 2019

PROTOCOL integer ID:
17662