VERSION 1

JAN 01, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.4r5gv86

External link:
<https://www.addgene.org/protocols/primer-design/>

Protocol Citation: Addgene The Nonprofit Plasmid Repository 2020. How to Design a Primer.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.4r5gv86>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Jun 26, 2019

Last Modified: Aug 17, 2020

How to Design a Primer V.1

In 2 collections

Addgene The Nonprofit Plasmid Repository¹

¹Addgene

Addgene The Nonprofit Plasmid Repository
 Addgene

ABSTRACT

This protocol describes how to design a primer. To see the full abstract and additional resources, visit <https://www.addgene.org/protocols/primer-design/>.

GUIDELINES

General Guidelines

The main property of primers is that they must correspond to sequences on the template molecule (must be complementary to template strand). However, primers do not need to correspond to the template strand completely; it is essential, however, that the 3' end of the primer corresponds completely to the template DNA strand so elongation can proceed. Usually a guanine or cytosine is used at the 3' end, and the 5' end of the primer usually has stretches of several nucleotides. Also, both of the 3' ends of the hybridized primers must point toward one another.

The size of the primer is very important as well. Short primers are mainly used for amplifying a small, simple fragment of DNA. On the other hand, a long primer is used to amplify a eukaryotic genomic DNA sample. However, a primer should not be too long (> 30-mer primers) or too short. Short primers produce inaccurate, nonspecific DNA amplification product, and long primers result in a slower hybridizing rate. On average, the DNA fragment that needs to be amplified should be within 1-10 kB in size.

The structure of the primer should be relatively simple and contain no internal secondary structure to avoid internal folding. One also needs to avoid primer-primer annealing which creates primer dimers and disrupts the amplification process. When designing, if unsure about what nucleotide to put at a certain position within the primer, one can include more than one nucleotide at that position termed a mixed site. One can also use a nucleotide-based molecular insert (inosine) instead of a regular nucleotide for broader pairing capabilities.

Taking into consideration the information above, primers should generally have the following properties:

- Length of 18-24 bases
- 40-60% G/C content
- Start and end with 1-2 G/C pairs
- Melting temperature (T_m) of 50°C | 60°C
- Primer pairs should have a T_m within 5°C of each other
- Primer pairs should not have complementary regions

Note

Note, if you will be including a restriction site at the 5' end of your primer, note that a 3-6 base pair "clamp" should be added upstream in order for the enzyme to cleave efficiently (e.g. GCGGCG-restriction site-your sequence).