VERSION 1

MAR 09, 2019

HPLC sample prep V.1

Kaitly Woodard¹

¹University of Tennessee Health Science Center

Kaitly Woodard

ABSTRACT

Preparation of blood for hemoglobin separation (ion-exchange and reverse-phase HPLC).

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.yy3fxyn

Protocol Citation: Kaitly Woodard 2019. HPLC sample prep. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.yy3fxyn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Mar 09, 2019

Last Modified: Mar 09, 2019

PROTOCOL integer ID:
21243

1 Prepare 1.5mL Eppendorf tubes with 200uL of Hemolysate Reagent

- 2 Add 1uL of blood to tube, and vortex for 5 seconds.
- 3 Centrifuge at max speed for 10 minutes
- 4 Pipette out 75-100uL of supernatant into HPLC vial, careful not to disturb DNA pellet at the bottom of the tube. If pipette tip appears sticky after removal from supernatant you may have taken the DNA. Instead, discard the sticky debris and pipette from remainder of supernatant. (The DNA and cell debris will stay together, once it is removed only desired hemolysate will remain).
- 5 Ensure vial is labeled appropriately, and no bubbles are present at the bottom of the vial. Bubbles can be removed by tapping the vial gently on the benchtop or removing by pipette.
- 6 **Reserve HPLC time on the calendar, provide sample list and samples to JJZ