AUG 14, 2020

Human Islet Microvasculature Immunofluorescence in Optically Cleared Samples

Martha Campbell-Thompson¹, Elizabeth Butterworth Hosaka², Katelyn N Carty²

¹University of Florida; ²co-author

Human BioMolecular Atlas Program (HuBMAP) Method Development Community

Tech. support email: Jeff.spraggins@vanderbilt.edu

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.y3tfynn

External link:
<https://www.protocols.io/view/human-pancreas-pact-optical-clearing-and-high-reso-9gbh3sn/materials>

Protocol Citation: Martha Campbell-Thompson, Elizabeth Butterworth Hosaka, Katelyn N Carty 2020. Human Islet Microvasculature Immunofluorescence in Optically Cleared Samples . **protocols.io** <https://dx.doi.org/10.17504/protocols.io.y3tfynn>

MANUSCRIPT CITATION: Butterworth E, Dickerson WD, Vijay V, Weitzel K, Cooper J, Atkinson EW, Coleman JE, Otto KJ, Campbell-Thompson M. High resolution 3D Imaging of the Human Pancreas Neuro-Insular Network. J Vis Exp. 2018 Jan 29;(131). doi: 10.3791/56859.2018. PMID: 29443037.

Jesus Peñaloza
University of Florida

DISCLAIMER

Different primary and secondary antibody lots may differ in affinities and should be independently optimized.

ABSTRACT

This protocol describes the immunostaining performed on five human control pancreas samples, cleared using a modified passive CLARITY (PACT) method according to our published methods. Islets were identified based on alpha-cells stained with glucagon while collagen-containing basement membranes in the extracellular matrix were stained using anti-collagen IV. Smooth muscle cells surrounding arteries and arterioles were identified using anti-SMA primary antibody directly conjugated with Cy3 fluorophore. An accompanying protocol describes using Vesselucida360 to contour islets and determine islet basement membrane and SMA densities and morphometric variables (<https://dx.doi.org/10.17504/protocols.io.bjfkj6>).

ATTACHMENTS

[jove-protocol-56859-high-resolution-3d-imaging-of-the-human-pancreas-neuro-insular-network.pdf](#)

IMAGE ATTRIBUTION

An islet is shown from case 6232 with all 3 fluorescent channels visible.

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Mar 12, 2019

Last Modified: Aug 14, 2020

PROTOCOL integer ID:
21331

Keywords: human islet, glucagon, optical clearing, confocal microscopy, 3D, PACT, collagen IV, SMA, Vesselucida360, alpha-cells, pancreas, neuro-insular network, passive CLARITY

GUIDELINES

This protocol is for PACT-cleared human pancreas samples, fixed with 4% paraformaldehyde for at minimum 48:00:00, 4% agarose embedded and sectioned at 400 um or greater thickness. The Triton-X concentration in the antibody diluent was increased from 0.1% to 0.5% based on experiments that showed antibody penetration into the middle of the sample was improved without loss of signal.

MATERIALS

MATERIALS

Glucagon (GCG) Mouse anti-human **Abcam Catalog #ab10988 (Lot# GR290488-2)**

Goat anti-ms Dylight 405 hi-cross **Thermo Fisher Scientific Catalog #35500BID**

PBS Phosphate Buffered Saline 10X Solution **Fisher Scientific Catalog #BP399-1**

Goat normal serum **Vector Laboratories Catalog #S-1000**

Triton X-100 **Merck MilliporeSigma (Sigma-Aldrich) Catalog #T8787-100**

Collagen IV (Col14) Rabbit anti-human **Abcam Catalog #ab6586**

Smooth muscle actin- alpha (a-SMA) Mouse anti-human-Cy3 [clone 1A4] **Sigma - Aldrich Catalog #C6198**

Goat anti-rb AF488 hi cross **Thermo Fisher Scientific Catalog #A-11034**

STEP MATERIALS

Triton X-100 **Merck MilliporeSigma (Sigma-Aldrich) Catalog #T8787-100**

PBS Phosphate Buffered Saline 10X Solution **Fisher Scientific Catalog #BP399-1**

Software

Vesselucida 360

NAME

MBF Bioscience

DEVELOPER

Software

Vesselucida Explorer

NAME

MBF Bioscience

DEVELOPER

SAFETY WARNINGS

Follow all laboratory safety procedures when handling hazardous chemicals and materials.

BEFORE START INSTRUCTIONS

Ensure samples are cleared and the SDS is well-removed by extensive washing in PBS at Room temperature on a rocker plate.

- 1 For these samples, optical clearing was performed using 4% SDS at 60 °C with rocking for up to 3 weeks until the samples reached transparency. Our original protocol used 8% SDS. We have subsequently tested both 4% and 8% SDS at Room temperature and 60 °C. Different tissues should be tested independently for optimal results.

Prepare the Antibodies

- 2 Make antibody dilution buffer (ABD): 1x PBS containing 2% normal goat serum (NGS) and 0.5% Triton-X 100.

PBS Phosphate Buffered Saline 10X Solution **Fisher Scientific Catalog #BP399-1**

Goat normal serum **Vector Laboratories Catalog #S-1000**

Triton X-100 **Merck MilliporeSigma (Sigma-Aldrich) Catalog #T8787-100**

- 3 Dilute the primary antibodies to 1:200 in ABD and prepare at least 1 mL per sample.

Stain the Tissue

- 4 Incubate the samples with primary antibodies on a rocker plate for 48:00:00 1-5 days at Room temperature .

- 5 Wash the tissue sections with 1xPBS for five times, minimum 01:00:00 at Room temperature .

PBS Phosphate Buffered Saline 10X Solution **Fisher Scientific Catalog #BP399-1**

- 6 Dilute the secondary antibodies to 1:500 and prepare at least 1 mL per sample. Incubate for 48:00:00 or more on a rocker at Room temperature .

- 7 Wash the tissue with 1 x PBS 5 times 01:00:00 at Room temperature .

- 8 Equilibrate the samples in RIMS containing 0.5% sodium azide for at least 12:00:00 before imaging.
Transfer to an imaging dish or chamber slide using RIMS as mounting media.

Imaging

- 9 Perform confocal microscopy. Each microscope will require optimization of imaging parameters.

Equipment

LSM 710

NAME

confocal laser microscope

TYPE

Zeiss

BRAND

LSM 710

SKU

<https://www.zeiss.com/microscopy/us/products/confocal-microscopes.html>

LINK

- 10 The following lasers were used for imaging each antigen in these images:

Track 1- 488- Collagen IV

Track 2- 405- GCG, 561- SMA

Results

11

Expected result

Islet alpha-cells- GCG Stain - Specific Stain, Intensity Good, Background Low

Collagen IV- basement membranes- Specific Stain, Intensity Good, Background medium

SMA- smooth muscle cells surrounding vessels - Specific Stain, Intensity Good, Background Low

- 12 Analyze islet microvascular density by contouring the GCG+ area for islet volume followed by collagen IV and SMA morphometry using Vesselucida360.