NOV 01, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.7pyhmpw

Protocol Citation: Paul Gerald Layague Sanchez, Hidenobu Miyazawa, Molecular Instruments 2019. Hybridization chain reaction (HCR) protocol for tails of mouse embryos. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.7pyhmpw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol in our group and it is working.

Created: Sep 26, 2019

Last Modified: Nov 01, 2019

PROTOCOL integer ID:
 28120

Keywords: HCR, in situ hybridization

Hybridization chain reaction (HCR) protocol for tails of mouse embryos

Paul Gerald Layague Sanchez¹, Hidenobu Miyazawa¹, Molecular Instruments²

¹European Molecular Biology Laboratory (EMBL); ²Molecular Instruments

Paul Gerald Layague Sanchez
 European Molecular Biology Laboratory (EMBL)

ABSTRACT

This protocol is for the in situ hybridization chain reaction (HCR) of intact tails (somites and presomitic mesoderm, PSM) of mouse embryos. This is adapted from the in situ HCR v 3.0 protocol for whole-mount mouse embryos available in the website of Molecular Instruments (for details, please follow this [link](#)).

References

- Choi, H., Schwarzkopf, M., Fornace, M., Acharya, A., Artavanis, G., & Stegmaier, J. et al. (2018). Third-generation in situ hybridization chain reaction: multiplexed, quantitative, sensitive, versatile, robust. *Development*, 145(12), dev165753. doi: [10.1242/dev.165753](https://doi.org/10.1242/dev.165753)
- Choi, H., Calvert, C., Husain, N., Huss, D., Barsi, J., & Deverman, B. et al. (2016). Mapping a multiplexed zoo of mRNA expression. *Development*, 143(19), 3632-3637. doi: [10.1242/dev.140137](https://doi.org/10.1242/dev.140137)

Day 1: Recovery and fixation of samples

- 1 Recover samples, wash quickly with cold PBS, and fix in 4% formaldehyde (1.08 mL 37% formaldehyde diluted to 10 mL with PBS). Keep in formaldehyde overnight (around 16:00:00) at 4 °C in the cold room, with gentle shaking/rolling.

Safety information

Formaldehyde is toxic. Operate under a fume hood and wear appropriate hand, body, and eye protection.

Note

To reduce autofluorescence, it is recommended to use fresh fixative and PBS with no divalent cations (like Ca^{2+} and Mg^{2+}).

Day 2: Dehydration of samples

- 2 Wash samples with PBST (0.1% Tween20 in 1x PBS).
 - Wash 1/2 00:05:00
 - Wash 2/2 00:05:00
- 3 Dehydrate samples (one sample per tube) with a series of graded methanol : PBST (PBS + 0.1% Tween) washes.
 - 25% methanol : 75% PBST 00:05:00
 - 50% methanol : 50% PBST 00:05:00
 - 75% methanol : 25% PBST 00:05:00
 - 100% methanol 00:05:00
 - 100% methanol 00:05:00
- 4 Incubate samples at -20 °C overnight (around 16:00:00) or until use.

Day 3: Rehydration of samples and HCR detection stage

- 5 Rehydrate samples (one sample per PCR tube, capacity = 200 μ L max) with a series of graded methanol : PBST (PBS + 0.1% Tween) washes.
- 100% methanol 00:05:00
 - 75% methanol : 25% PBST 00:05:00
 - 50% methanol : 50% PBST 00:05:00
 - 25% methanol : 75% PBST 00:05:00
 - 100% PBST 00:05:00
 - 100% PBST 00:05:00
- 6 Treat samples with 10 μ g/mL proteinase K (Merck, CAS # 38450-01-6) for 00:05:00 at Room temperature .
- 7 Wash samples with PBST.
- Wash 1/2 00:05:00
 - Wash 2/2 00:05:00
- 8 Fix samples in 4% formaldehyde (2.16 mL 37% formaldehyde diluted to 20 mL with PBST) for 00:20:00 at Room temperature .

Pre-warm probe hybridization (PH) buffer at 37 $^{\circ}$ C for later use.

Safety information

Formaldehyde is toxic. Operate under a fume hood and wear appropriate hand, body, and eye protection.

Safety information

Probe hybridization (PH) buffer contains formamide, which is toxic.

Note

To reduce autofluorescence, it is recommended to use fresh fixative and PBS with no divalent cations (like Ca^{2+} and Mg^{2+}).

9 Wash samples with PBST.

- Wash 1/3 00:05:00
- Wash 2/3 00:05:00
- Wash 3/3 00:05:00

10 Wash samples with PH buffer for 00:05:00 .

Safety information

Probe hybridization (PH) buffer contains formamide, which is toxic.

11 Put fresh PH buffer to the tubes and incubate for 00:30:00 at 37 °C .

Meanwhile, prepare probe solution by adding 2 pmol (2 μL of [M] 1 micromolar (μM) stock) of odd probe mixture and 2 pmol of even probe mixture to every 500 μL of pre-warmed PHP buffer (from Step 8).

Safety information

Probe hybridization (PH) buffer contains formamide, which is toxic.

12 Remove PH buffer and add the probe solution. Incubate overnight (around 16:00:00) at 37 °C .

Day 4: HCR amplification stage

13 Pre-warm probe wash buffer at 37 °C , and equilibrate amplification buffer to Room temperature .

Safety information

Probe wash (PH) buffer contains formamide, which is toxic.

- 14** In separate tubes, for every 500 μL hairpin mixture, prepare 30 pmol (15 μL of 2 micromolar (μM) stock or 10 μL of 3 micromolar (μM) stock) of hairpin h1 (tube 1) and 30 pmol of hairpin h2 (tube 2). Snap-cool hairpins by heating the tubes at 95 °C for 00:01:30 and cool to Room temperature in the dark for at least 00:30:00.

- 15** Wash samples with probe wash buffer at 37 °C.

- Wash 1/4 00:15:00
- Wash 2/4 00:15:00
- Wash 3/4 00:15:00
- Wash 4/4 00:15:00

Safety information

Probe wash (PH) buffer contains formamide, which is toxic.

- 16** Wash samples with 5x SSCT (0.1% Tween20) at Room temperature.

- Wash 1/2 00:05:00
- Wash 2/2 00:05:00

Note

To prepare 500 mL of 5x SSCT, dilute 125 mL of 20x SSC and 0.5 mL Tween20 to 500 mL with distilled H₂O.

- 17** Wash samples with pre-equilibrated amplification buffer (Step 13) for 00:05:00 at Room temperature.

Meanwhile, prepare the hairpin mixture (30 pmol of each hairpin in 500 μL mixture) by mixing the snap-cooled hairpins h1 and h2 (Step 14) to amplification buffer at

Room temperature

- 18 Remove the amplification buffer from the sample tubes, and add the hairpin mixture. Incubate overnight (around 16:00:00) in the dark at Room temperature .

Note

For dHCR imaging, amplify for a shorter period of time to ensure single-molecule dots are diffraction-limited.

Day 5: Washing, nuclear staining, and imaging

- 19 Wash samples with 5x SSCT for 00:30:00 in the dark at Room temperature .

- 20 Stain nuclei with 1:1000 DAPI (5 ug/mL : 20 μ L of 5 mg/mL DAPI diluted to 20 mL with 5x SSCT) in the dark at Room temperature .

- Wash + DAPI 1/3 00:30:00
- Wash + DAPI 2/3 00:30:00
- Wash + DAPI 3/3 00:30:00

- 21 Wash samples with 5x SSCT in the dark at Room temperature

- Wash 1/2 00:30:00
- Wash 2/2 00:30:00

- 22 Samples are now ready for imaging.

Prior to imaging, the samples could be cleared using the fructose-glycerol clearing solution described in [Dekkers et al., 2019](https://doi.org/10.1016/j.cub.2019.05.001). Incubate samples in clearing solution in the dark at 4 °C for at least 02:00:00 .

Note

To prepare the fructose-glycerol clearing solution, dissolve 29.72 g fructose in 33 mL glycerol and 7 mL water on a magnetic stirrer (start the dissolution at least a day before intended day of clearing because it will take a while for fructose to dissolve). The clearing solution can be stored in the dark at 4 °C for at most 1 month. This solution can be used to mount the sample on microscope slide/cover glass. Take note that the clearing is solution has high viscosity.

Expected result

Tbx18 and Uncx marking the two halves (anterior and posterior, respectively) of somites in tail of a mouse embryo. Transcripts were visualized using HCR version 3.0. Nuclei were stained using DAPI.