MAR 06, 2020

Hydroxyl Radical-scavenging Activity

PLOS One

Jing Xu¹

¹Hainan University

Jing Xu

Hainan University

ABSTRACT

The scavenging of hydroxyl radicals were determined by Guo et al. The reactions were performed in 0.3 ml of 20 mM sodium salicylate, 2.0 ml of 1.5 nM FeSO₄, 1.0 ml of samples, and 1.0 ml of 6 mM H₂O₂. The reaction mixture was incubated for 1 hat 37 °C. The absorbance was measured at 510 nm. The inhibition percent and 50% of absorbance(EC₅₀)were calculated.

OPEN ACCESS

DOI:

[dx.doi.org/10.17504/protocols.io.bdazi2f6](https://doi.org/10.17504/protocols.io.bdazi2f6)

External link:

<https://doi.org/10.1371/journal.pone.0234435>

Protocol Citation:

Jing Xu
2020. Hydroxyl Radical-scavenging Activity.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.bdazi2f6>

MANUSCRIPT CITATION:

Zhou J, Yang Q, Zhu X, Lin T, Hao D, Xu J (2020) Antioxidant activities of *Clerodendrum cyrtophyllum* Turcz leaf extracts and their major components. PLoS ONE 15(6): e0234435. doi: [10.1371/journal.pone.0234435](https://doi.org/10.1371/journal.pone.0234435)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status:

Working
We use this protocol and it's working

Created: Mar 06, 2020

Last Modified: Mar 06, 2020

PROTOCOL integer ID:
33849