DEC 07, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.984h9yw

External link:
<https://doi.org/10.1371/journal.pone.0228823>

Protocol Citation: Ding Mingliang, Muhammad Asim, Li Mingju, Sedhom Abdelkhalik, Daniel Manore, Li Shaoxiang, Zhao Hong, Lin Liping 2019. Identification of promising genotypes. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.984h9yw>

MANUSCRIPT CITATION:
 Mingliang D, Asim M, Mingju L, Abdelkhalik S, Manore D, Shaoxiang L, Hong Z, Liping L (2020) Wheat genetic gains for two distinct management schemes in China: An analysis of elite spring type genotypes. PLoS ONE 15(2): e0228823. doi: [10.1371/journal.pone.0228823](https://doi.org/10.1371/journal.pone.0228823)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Identification of promising genotypes

PLOS One

Ding Mingliang¹, Muhammad Asim^{1,2}, Li Mingju³, Sedhom Abdelkhalik^{1,4}, Daniel Manore^{1,5}, Li Shaoxiang¹, Zhao Hong¹, Lin Liping⁶

¹Institute of Food Crops, Yunnan Academy of Agricultural Sciences, Kunming, China;

²Pakistan Agricultural Research Council, Islamabad, Pakistan;

³Institute of Agricultural Environment and Resources, Yunnan Academy of Agricultural Sciences, Kunming, China;

⁴Field Crops Research Institute, Agricultural Research Center, Egypt;

⁵Department of Plant Science, College of Agriculture, Wachemo University, Ethiopia;

⁶Seed Management Department of Yunnan Province, Kunming, China

Muhammad Asim

Protocol status: Working

We use this protocol in our group and demonstrated successful identification of promising genotypes for field as well as upland wheat management schemes in China

Created: Dec 07, 2019

Last Modified: Dec 07, 2019

PROTOCOL integer ID:

30716

- 1 Average the yield data generated for genotypes across locations and years.
- 2 Perform the Welch's variance-weighted analysis of variance for studied genotypes (across years and locations), at each location (across genotypes and years) and for each year (across genotypes and locations) using One-Way ANOVA model of SAS University Edition version 3.8 with Levene test for homogeneity of variance and adopting Tukey comparison method.
- 3 Calculate mean genetic deviation in successive cycles
- 4 Corresponding above mean values facilitate the partitioning of promising genotypes, location and years