NOV 15, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.8rmhv46

Protocol Citation: Elizabeth Smith 2019. IHC-Amplified Fluorescent Frozen Sections. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.8rmhv46>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Oct 25, 2019

Last Modified: Nov 15, 2019

PROTOCOL integer ID:
29197

IHC-Amplified Fluorescent Frozen Sections

Elizabeth
Smith¹

¹East Tennessee State University

SPARC

Tech. support email: info@neuinfo.org

Elizabeth Smith

ABSTRACT

Immunohistochemistry protocol used for staining with fluorescent secondary antibodies to highlight specific tissue structures and amplify specific antibody signals.

ATTACHMENTS

[Fluor Immun Protocol-Biotin_Streptavidin-Slides \(amplification\).docx](#)

GUIDELINES

Frozen tissues cut in 30 micron intervals were used in this protocol. Do not touch the tissue on the slide or it will come off.

MATERIALS

MATERIALS

- ✕ Normal Donkey Serum Contributed by users Catalog #017-000-121
- ✕ Citifluor AF-1 anti-fading solution Electron Microscopy Sciences
- ✕ KimWipes Fischer Scientific
- ✕ Triton X-100 Sigma Catalog #93426
- ✕ PBS Contributed by users
- ✕ ImmEdge hydrophobic barrier pap pen Vector Laboratories Catalog #H-4000
- ✕ Superfrost™ Disposable Microscope Slides, White; 3 x 1 in. x 1mm Thermo Fisher Catalog #12550123
- ✕ Bovine Serum Albumin Thermo Fisher Catalog #15561020
- ✕ Coplin Staining Jar Thermo Fisher Catalog #194
- ✕ Scientific Device Humidity/Slide Moisture Chamber Thermo Fisher Catalog #23769522

Slides boxes wrapped in tinfoil to store slides in -20 degrees Celsius prior to cutting.

Various primaries and secondaries dependent upon structure of interest.

Glass coverslips in various sizes depending on tissue size.

Primaries:

Neuropeptide Y (NPY) Antibody from Immunostar Catalog#22940

Anti-Tyrosine Hydroxylase (TH) from Millipore Catalog#AB1542

VACHT from Synaptic Systems Catalog#139 103

Anti-PGP9.5 antibody from abcam Catalog#ab108986

Secondaries:

Alexa Fluor 488 conjugated AffiniPure Donkey Anti-Rabbit IgG from Jackson ImmunoResearch Catalog#711-545-152

Alexa Fluor 555 conjugated Donkey Anti-Goat IgG from ThermoFisher Catalog#A-21432

Alexa Fluor 555 conjugated Donkey Anti-Rabbit IgG from ThermoFisher Catalog#A-31572

Alexa Fluor 594 conjugated Donkey Anti-Rabbit IgG from Jackson ImmunoResearch Catalog#711-585-152

Alexa Fluor 594 conjugated Donkey Anti-Goat IgG from ThermoFisher Catalog#A-11058

Biotin-SP-conjugated Affinipure Donkey Anti-Goat IgG from Jackson ImmunoResearch Catalog#705-065-147

Streptavidin from Thermo Scientific Catalog#21842

- 1 Day 1: Using the PAP Pen, carefully draw a water barrier circle around the tissue sections on the slide – allow this circle to dry for several seconds or up to approx. one minute
- 2 Rinse slides with PBS (pH 7.3-7.4): 4 x 5 min each
- 3 Rinse slides with 0.5% BSA + 0.4% Triton X-100 in PBS): 1 x 10 min
- 4 Remove slides one at a time and using a clean Kimwipe, carefully wipe around the tissue sections to dry the slide
- 5 Place the slides into a black, covered slide incubation box/humidity box
- 6 Cover the tissue sections with blocking buffer (10% normal donkey serum in 1.0% BSA + 0.4% Triton X-100 + PBS)
- 7 Allow the sections to remain in blocking buffer for 1.5-2 hrs. at RT
- 8 Pour off the blocking buffer
- 9 Replace with primary antibody solution (antibody of choice diluted in 1.0% BSA + 0.4% Triton X-100 + PBS)

- 10** Incubate tissue with primary antibody overnight in incubation box
- 11** Day 2: Rinse slides with PBS: 4 x 5 min
- 12** Rinse slides with 0.5% BSA + 0.4% Triton X-100 + PBS: 1 x 10 min
- 13** Place the slides into a black, covered slide incubation box/humidity box
- 14** Cover the tissue sections with blocking buffer (10% normal donkey serum in 1.0% BSA + 0.4% Triton X-100 + PBS)
- 15** Allow the sections to remain in blocking buffer for 1.5-2 hrs. at RT
- 16** Prepare fluorescent secondary antibody (secondary antibody should be diluted in 1.0% BSA + 0.4% Triton X-100 + PBS)
- 17** Cover the tissue with the secondary antibody solution and incubate for 2 hrs. at RT in the incubation box. *From this point on, use low light and/or cover tissues.*

- 18 Rinse slides with PBS: 4 x 5 min
- 19 Remove excess PBS with a Kimwipe
- 20 Prepare fluorescent Streptavidin secondary antibody (should be diluted in 1.0% BSA + 0.4% Triton X-100 + PBS)
- 21 Cover the tissue with the fluorescent Streptavidin antibody solution and incubate for 2 hrs. at RT in the incubation box.
- 22 Rinse slides with PBS: 4 x 5 min
- 23 Remove excess PBS with a Kimwipe
- 24 Carefully add a drop of mounting medium to the center of the tissue and apply cover glass
- 25 Seal cover glass with clear nail polish. For thicker tissue, add a weight before sealing