MAY 13, 2018

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.p4hdqt6

External link:
<https://doi.org/10.1371/journal.pone.0201453>

Protocol Citation: Souza TKF, Nucci MP, Mamani JB, Silva HR, Fantacini DMC, Souza LEB, Picanço-Castro V, Covas DT, Vidoto EL, Tannús A, Gamarra LF 2018. Image and behavior analysis in C6 glioma model. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.p4hdqt6>

MANUSCRIPT CITATION: Souza TKF, Nucci MP, Mamani JB, Silva HRd, Fantacini DMC, Souza LEBd, Picanço-Castro V, Covas DT, Vidoto EL, Tannús A, Gamarra LF (2018) Image and motor behavior for monitoring tumor growth in C6 glioma model. PLoS ONE 13(7): e0201453. doi: [10.1371/journal.pone.0201453](https://doi.org/10.1371/journal.pone.0201453)

Image and behavior analysis in C6 glioma model

PLOS One

Souza TKF¹, Nucci MP², Mamani JB¹, Silva HR¹, Fantacini DMC³, Souza LEB³, Picanço-Castro V³, Covas DT³, Vidoto EL⁴, Tannús A⁴, Gamarra LF¹

¹Hospital Israelita Albert Einstein, São Paulo, SP, Brazil;

²LIM44, Faculdade de Medicina, Universidade de São Paulo, São Paulo, SP - Brazil.;

³Faculdade de Medicina de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, SP - Brazil;

⁴CIERMag-Instituto de Física de São Carlos, Universidade de São Paulo, São Paulo, Brazil

Nanobiotechnolgy

Lionel Gamarra

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: May 13, 2018

Last Modified: May 13, 2018

PROTOCOL integer ID:
12137

Keywords: glioblastoma, gait analysis, behavioral, CatWalk, motor function, Bioluminescence, MRI, Actimeter, glioma, tumor

Tumor Induction

- 1
 - The animals are weighed and anesthetized with an *i.p.* injection of ketamine (100 mg/kg) and xylazine (20 mg/kg) and place in a stereotaxic apparatus.
 - The C6 (10^6 cells/30 mL) cells are implanted. The implantation position is determined at the following coordinates: 2.0 mm anteroposterior, 2.0 mm laterolateral, and a depth of 2.5 mm. The cells are diluted to a concentration of 10^6 cells/30 mL.
 - The cells are slowly injected over a 10-min period. For the control group, culture medium is injected without the cells.
 - The bone is then reassembled by using bone wax, and the skin is sutured using cotton thread.

Tumor growth monitoring by MRI and bioluminescence imag.

- 2
 - The animals are weighed and anesthetized with an *i.p.* injection of ketamine (100 mg/kg) and xylazine (20 mg/kg)
 - The MRI and bioluminescence images are acquired at base time and at 7, 14, 21, and 28 days after tumor induction surgery for tumor growth analysis.
 - T2-weighted images are acquired by using a rapid acquisition with relaxation enhancement sequence (RARE).
 - The total tumor volume is calculated by the sum of tumor area in each slice multiplied by the thickness and gap per slice.
 - In each *in vivo* bioluminescence imaging procedure, rats are anesthetized with isoflurane and

D-Luciferin is injected i.p. at a dose of 450 mg/kg 10 min before of the BLI acquirement in IVIS.

- Afterward, for *ex vivo* BLI acquiring, one animal per group of the animals that acquired the *in vivo* BLI is euthanized, and the same brain of *ex vivo* BLI is used for histological assessment.

Behavioral : testing Spontaneous locomotor activity

- 3
 - Spontaneous global locomotor activity are quantified by the Infrared (IR) Actimeter. The apparatus comprises a two-dimensional (X-axis and Y-axis) square frame of 450 × 450 mm², surrounded by transparent walls of 30 cm high, a frame support, and a control unit.
 - Each animal are placed in the center of the arena, and the spontaneous locomotor behavior is tracked for 5 min.
 - The six parameters used for comparison between groups and sessions are slow movements (S-MOV), fast movements (F-MOV), slow stereotyped (S-STE), fast stereotyped (F-STE) slow rearing (S-REA), and fast rearing (F-REA).

Behavioral : Gait assessment by CatWalk

- 4
 - The gait is analyzed by using a "Catwalk" (Noldus Information Technology, Netherlands) apparatus that comprises a 1.3-m long glass platform illuminated by fluorescent lights that are reflected downward while pressure is applied from above.
 - During data analysis, each the print on paw contact is automatically classified as right forepaw (RF), right hindpaw (RH), left forepaw (LF), and left hindpaw (LH).
 - After the identification of individual footprints, we performed an automated analysis of wide-range parameters. Data are classified as follows:
 - **Spatial parameters:** print positions (PP), the distance between the placement of a hindpaw and the ipsilateral frontpaw placed just before it; base of support (BOS), the distance between the center points of the two fore or hind paws (both represent inter-limb coordination measures); stride length (SL), the distance between successive placements of the same paw (dynamic gait parameters); and maximum contact area (mCA), the maximum area of a paw that comes into contact with the glass plate (static gait parameters).
 - **Temporal parameters:** **stand**, the duration in seconds of contact of a paw with the glass plate; **step cycle (SC)**, the duration of stance and swing phases combined; **cadence**, steps per second; and **duration**, total time of entire run.