



Aug 14, 2020

Immunoblot analyses for investigating SpLA binding to purified mammalian and avian immunoglobulins.

DOI

dx.doi.org/10.17504/protocols.io.bjq7kmzn

Angel A Justiz-Vaillant¹

¹University of the West Indies St. Augustine

University of the West In...

angel.vaillant@sta.uwi.e...



Angel A Justiz-Vaillant

University of the West Indies St. Augustine

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.bjq7kmzn

Protocol Citation: Angel A Justiz-Vaillant 2020. Immunoblot analyses for investigating SpLA binding to purified mammalian and avian immunoglobulins.. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bjq7kmzn>

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: August 14, 2020

Last Modified: August 14, 2020

Protocol Integer ID: 40447

- 1 Aliquots of 2 $\mu\text{g}/\mu\text{l}$ purified immunoglobulins from birds, laboratory, wild, farm animals and pets are applied to the gels of SDS-PAGE as described elsewhere.
- 2 Gels are transferred to nitrocellulose membranes (Immobilon-Nc, pore size 0.45 μm , Sigma-Aldrich Co, St Louis, Missouri) during 75 minutes at 40 mAmps using a semi-dry electroblotter, HEP-1 Model, Owl Scientific Inc.
- 3 The running buffer contains 25 mM Tris, 192 mM glycine pH 8.3 and 20% methanol.
- 4 The nitrocellulose membranes are blocked overnight in 10% non-fat skim milk in PBS with 0.05% Tween-20 pH 7.4 and then washed 4x, 10 minutes with PBS-Tween 20.
- 5 Recombinant protein LA (Sigma-Aldrich Co, St Louis, Missouri) at a concentration of 5 $\mu\text{g}/\text{ml}$ is added to the membranes.
- 6 After that there is an incubation period at 4°C overnight.
- 7 The nitrocellulose membranes were washed as above.
- 8 A secondary antibody (rabbit anti-chicken IgY horseradish peroxidase, Sigma Aldrich) is added at a 1:15 000 dilution.
- 9 It is incubated for one hour at room temperature and washed as above.
- 10 Tetramethyl-benzidine (TMB) solution is added to the nitrocellulose membranes, which are then incubated in the dark for a few minutes. Then, the membranes are shaken gently and rinsed thoroughly in de-ionized water to stop the blotting process and are left to dry.
- 11 Alternatively, Ig samples are transferred to nitrocellulose membranes and directly probed using SpLA-HRP (diluted 1:5000) and TMB (this system was mainly used for detecting avian Igs).