



Jul 10, 2024

Immunocytochemistry

 The Journal of Cell Biology

DOI

<https://dx.doi.org/10.17504/protocols.io.14egn2556g5d/v1>

Shiyi Wang¹

¹Duke University



Shiyi Wang

Duke University

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.14egn2556g5d/v1>

External link: <https://doi.org/10.1083/jcb.202410130>

Document Citation: Shiyi Wang 2024. Immunocytochemistry. **protocols.io**

<https://dx.doi.org/10.17504/protocols.io.14egn2556g5d/v1>

Manuscript citation:

Salazar MPR, Kolanukuduru S, Ramirez V, Lyu B, Manigault G, Sejourne G, Sesaki H, Yu G, Eroglu C (2025) Mitochondrial fission controls astrocyte morphogenesis and organization in the cortex. The Journal of Cell Biology 224(10). doi: [10.1083/jcb.202410130](https://doi.org/10.1083/jcb.202410130)

License: This is an open access document distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Created: May 23, 2023

Last Modified: July 10, 2024

Document Integer ID: 82327

Keywords: ASAPCRN

Funders Acknowledgements:

Aligning Science Across Parkinson's (ASAP) initiative

Grant ID: ASAP-020607

Disclaimer

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](#) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](#), can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

Abstract

Immunocytochemistry

1. Astrocyte-neuron co-cultures on glass coverslips were fixed on DIV 12 with warm 4% PFA for 7 minutes, washed 3 times with PBS, blocked in a blocking buffer containing 50% normal goat serum (NGS) and 0.4% Triton X-100 for 30 minutes, and washed in PBS.
2. Samples were then incubated overnight at 4°C in primary antibodies diluted in blocking buffer containing 10% NGS, washed with PBS, incubated in Alexa Fluor conjugated secondary antibodies (Life Technologies) for 2 hours at room temperature, and washed again in PBS.
3. Coverslips were mounted onto glass slides (VWR Scientific) with Vectashield mounting media containing DAPI (Vector Labs), sealed with nail polish, and imaged on an Axiolmager M1 (Zeiss) fluorescence microscope.
4. Images of healthy astrocytes with strong expression of fluorescent markers that did not overlap with other fluorescent astrocytes were acquired at 40× magnification in red, green, and/or DAPI channels using a CCD camera.
5. Astrocyte morphological complexity was analyzed in FIJI using the Sholl analysis plugin, as described previously(<https://github.com/Eroglu-Lab/In-Vitro-Sholl>).
6. Statistical analyses were conducted using a custom code in R (<https://github.com/Eroglu-Lab/In-Vitro-Sholl>).
7. A mixed-effect model with multiple comparisons made using the Tukey post-test was used for Sholl analysis to account for the variability per experiment as a random effect to evaluate differences between the conditions.
8. The exact number of independent experiments and the exact number of cells analyzed are indicated in the figure legend for each experiment.
9. To ensure the health of astrocyte-neuron co-cultures, the peak for the number of astrocyte intersections must be greater than or equal to 25 in the control condition.
10. We imaged non-overlapping astrocytes that contained a single nucleus (DAPI stain) and expressed consistent fluorescent markers and EZRIN constructs according to the experimental conditions.