DEC 18, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.5wtg7en

External link:
<https://doi.org/10.1002/cpcb.51>

Protocol Citation: Michael S. Fernandopulle, Ryan Prestil, Christopher Grunseich, Chao Wang, Li Gan, Michael E. Ward 2019. IMMUNOCYTOCHEMISTRY OF i3NEURONS (Support Protocol 2). **protocols.io** <https://dx.doi.org/10.17504/protocols.io.5wtg7en>

MANUSCRIPT CITATION: Fernandopulle, M. S., Prestil, R., Grunseich, C., Wang, C., Gan, L., & Ward, M. E. (2018). Transcription-factor mediated differentiation of human iPSCs into neurons. Current Protocols in Cell Biology, e51. doi:<https://doi.org/10.1002/cpcb.51>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

IMMUNOCYTOCHEMISTRY OF i³NEURONS (Support Protocol 2)

In 1 collection

Michael S. Fernandopulle¹, Ryan Prestil¹, Christopher Grunseich¹, Chao Wang², Li Gan², Michael E. Ward¹

¹National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, Maryland;

²Gladstone Institute of Neurological Disease, Gladstone Institutes, San Francisco, California

Neurodegeneration Method Development Community

Tech. support email: ndcn-help@chanzuckerberg.com

Julia Rossmanith
 protocols.io

ABSTRACT

Staining i³Neurons for immunofluorescence (IF) studies is difficult due to the delicate and interconnected nature of the neuronal processes. These processes are easily disrupted in the extensive series of washes in an IF study, and initial dissociation of even a few processes often results in entire sheets of neurons lifting off the culture surface. The best ways to minimize these events are by reducing the number of total washes and by carrying out washes slowly on a tilted dish. Additionally, if possible, IF studies should only be done in neurons 10 days old or younger. After this point, neurons tend to be very delicate and even extremely gentle washing typically causes substantial cell washout.

ATTACHMENTS

[fernandopulle2018.pdf](#)

MATERIALS

- Fresh 16% paraformaldehyde (PFA) solution (Electron Microscopy Sciences, cat. no. 15710)

Paraformaldehyde, 16% (wt/vol) **Electron Microscopy Sciences Catalog #15710**

- Phosphate-buffered saline (PBS; e.g., Gibco, cat. no. 10010049)

Phosphate-buffered saline (PBS) without calcium or magnesium **Gibco - ThermoFisher Catalog #10010049**

Protocol status: Working
We use this protocol and it's working

Created: Jul 26, 2019

Last Modified: Dec 18, 2019

PROTOCOL integer ID:
26291

Keywords: i3LMN,
i3Neurons, iPSC, iPSC-derived
neurons, transcription factor-
mediated differentiation

- 10% (w/v) saponin solution (Acros Organics, cat. no. 419231000)

Saponin, From Quillaja Saponaria Molina Pract. **Fisher Scientific Catalog**
#AC419231000

- Donkey serum (Sigma, cat. no. D9663))

Donkey serum **Sigma Aldrich Catalog #D9663**

- 96-well plate with neurons to be fixed and stained (see protocols above)
- Antibodies (primary and secondary)
- Sodium azide (Sigma, cat. no. S8032)

Sodium azide **Sigma Aldrich Catalog #S8032**

- Hoechst/DAPI dye (20 mg/ml; 10,000×)
- Sterile filters (SteriFlip, Millipore)
- Liquid reagent reservoirs (Thermo, cat. no. 8096-11)

Equipment

Reagent Reservoirs

NAME

Reagent Reservoirs

TYPE

Thermo Scientific™

BRAND

8096-11

SKU

<https://www.thermofisher.com/order/catalog/product/95128095>

LINK

- P200 8-channel pipet (for processing 96-well dishes)
- Plate rocker
- Imaging system

SAFETY WARNINGS

Please see SDS (Safety Data Sheet) for hazards and safety warnings.

BEFORE START INSTRUCTIONS

NOTE: The following protocol assumes that the experimenter is using a full 96-well dish with the recommended cell counts and medium volumes as indicated in [Basic Protocols 6](#) and [8](#). Staining on other surfaces (i.e., 8-well chamber slides) may be performed by scaling volumes appropriately.

1 Make 15 mL of 8 % PFA solution (7.5 mL PBS, 7.5 mL 16 % PFA).

2 Make 50 mL of antibody blocking solution (3 % donkey serum and 0.1 % saponin in PBS). Filter sterilize.

Note

Detergents other than saponin (i.e., Triton X-100 or Tween-20 at 0.25 %) may also be used. Concentrations should be optimized by the user.

3 Pour the 8 % PFA solution into a liquid reagent reservoir.

4 Retrieve 96-well plate (45,000 to 50,000 cells/100 μ L medium in each well) and slowly add 100 μ L 8 % PFA solution to each well with 8-channel pipet. **Do not pipet up and down to mix.**

Note

4 % PFA is a typical fixative for ICC applications. This concentration is used here, but an aspiration step is eliminated by adding an equal volume of 8 %PFA directly to the culture medium on the cells. Pipetting up and down eliminates this advantage by promoting cell dissociation from the culture surface.

5 Incubate at Room temperature for 00:10:00 .

Note

Longer fixation times and/or cold incubations may be used as per requirements for particular antibodies.

- 6 Tilt dish to one side and lower 8-channel micropipet tips so that they contact the wall of each well. Slowly aspirate the PFA, leaving a small amount (if necessary) at the wall-culture surface interface.

Note

To prevent drying, aspirate only one column of wells at a time. Dispense PFA into a waste liquid reagent reservoir.

- 7 With plate tilted, slowly dispense 200 µL PBS onto the same wall in each well, taking care to direct the micropipet tip toward the wall and NOT the culture surface. Liquid should flow smoothly onto culture surface.

Note

PBS without detergent tends to adhere to the wall and then suddenly rush onto the culture surface all at once, promoting cell dissociation. To ensure more gradual flow, gently rub the micropipet tip against the wall in a side-to-side motion while dispensing. This action disrupts the surface tension of the dispensed PBS droplet, providing a mechanical substitute for detergent.

- 8 Repeat steps 6 and 7 for each column of wells.

[go to step #6](#)

- 9 Once fixative solution has been replaced with PBS, gently rock for 00:05:00 .

- 10 Repeat steps 6 to 9 two times, each with 100 µL washes of PBS.

go to step #6

- 11 Aspirate PBS (using plate tilt method).
- 12 Add 100 μ L of blocking solution to each well.
- 13 Gently rock at Room temperature for 00:30:00 .
- 14 During blocking, make up primary antibody solution(s) in blocking buffer or 3 % BSA solution.
- 15 Aspirate blocking solution.
- 16 Add primary antibody solution to plate (following procedure in steps 6 and 7).
- 17 Gently rock at Room temperature for 01:00:00 or at 4 °C overnight.
- 18 Aspirate primary antibody solution.

Note

Primary antibody may be saved and re-used for up to 1 month. Sodium azide (0.02 % final concentration) should be added to any saved antibody solutions to prevent microbial growth.

19 Add 150 μL to 200 μL of blocking solution to wells (following procedure in steps 6 and 7).

20 Gently rock for 00:05:00 .

21 Repeat steps 18 to 20 two times for a total of three washes.

go to step #18

22 Aspirate blocking solution.

23 Add secondary antibody solution (following procedure in steps 6 and 7).

24 Gently rock at Room temperature for 01:00:00 .

25 Wash wells with PBS three times (following procedure in steps 6 and 7).

Note

The second-to-last wash can contain Hoechst/DAPI dye if nuclear visualization is desired.

26 Image cells in PBS.

Note

Replace with mounting medium if desired.

27 Store plate at 4 °C .