APR 23, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bakcicsw

Protocol Citation: Janet R Keast, Peregrine B Osborne 2020. Immunohistochemical labeling of thick cryosections from pelvic ganglia.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.bakcicsw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Dec 16, 2019

Last Modified: Apr 23, 2020

PROTOCOL integer ID:
 31076

Keywords:
 immunohistochemistry;
 immunofluorescence

Immunohistochemical labeling of thick cryosections from pelvic ganglia

Forked from [Immunohistochemical analysis of ganglion neurons innervating the lower urinary tract \[keast-001-stage03\]](#)

In 1 collection

Janet R Keast¹, Peregrine B Osborne¹

¹University of Melbourne

SPARC

Tech. support email: info@neuinfo.org

Janet R Keast
 University of Melbourne

ABSTRACT

This protocol describes immunohistochemical procedures applied to thick (50 µm) cryosections mounted directly on slides. It is used when the structures to be analysed are too large to remain intact within thin (10-20 µm) cryosections. Antibodies have been selected to distinguish different neurochemical classes of autonomic ganglion neurons and synaptic boutons associated with these neurons. The protocol can also be used to characterize neurons containing retrograde tracer.

MATERIALS

MATERIALS

- ⊗ Horse serum **Sigma Aldrich Catalog #12449C**
- ⊗ OCT (Optimal Cutting Temperature compound) **Sakura Finetek Catalog #458**
- ⊗ Sheep anti-neuronal nitric oxide synthase antibody; AB_90743 **Merck Millipore Catalog #AB1529**
- ⊗ Cy3 Donkey anti-goat IgG **Jackson ImmunoResearch Catalog #705-165-147**
- ⊗ AF488 Donkey anti-rabbit IgG **Jackson ImmunoResearch Catalog #711-545-152**
- ⊗ PAP hydrophobic barrier pen **Enzo Life Sciences Catalog #ADI-950-233-0001**
- ⊗ Triton X-100 **Merck Millipore Catalog #X100**
- ⊗ Vectorshield anti-fade mounting medium **Vector Laboratories Catalog #H-1000**
- ⊗ Mouse anti-synaptophysin antibody **Agilent Technologies Catalog #M0776**
- ⊗ Rabbit anti-protein gene product 9.5 antibody **Merck Millipore Catalog #B592**

☒ Sheep anti-tyrosine hydroxylase antibody Merck Millipore Catalog #AB1542

☒ AF647 Donkey anti-mouse IgG Thermofisher Catalog #A-31571

Solutions:

- PBS: phosphate-buffered saline, 0.1 M, pH 7.2
- PBS containing 30% sucrose (w/v)
- PBS containing 0.1% sodium azide
- Blocking solution: PBS containing 10% non-immune horse serum and 0.5% triton X-100

Primary antibodies:

Abbreviation	Gene name	Synonym	RRID	Host species	Dilution
nNOS	Nos1	Neuronal nitric oxide synthase	AB_90743	sheep	1:2000
PGP	Uchl1	Protein gene product 9.5; ubiquitin C-terminal hydrolase 1	AB_11214054	rabbit	1:2000
SYP	Syp	Synaptophysin	AB_2199013	mouse	1:300
TH	Th	Tyrosine hydroxylase	AB_90755	sheep	1:1000

Neuronal cell bodies are labelled with PGP, nNOS or TH antibodies. Synaptic boutons associated with these cells are identified with synaptophysin-immunoreactivity.

Secondary antibodies:

Tag-antibody	Host species	Dilution
AF488 anti-rabbit	Donkey	1:1000
AF647 anti-mouse	Donkey	1:500
Cy3 anti-goat	Donkey	1:1000

Preparation of cryosections

- 1 Cryoprotect fixed tissue in phosphate-buffered saline (PBS; 0.1 M, pH7.2) containing 30% sucrose. This should be performed at 4°C, 24-72h prior to cutting.

- 2 Embed tissue in cryomold using OCT, freeze in cryostat and cut sections (50 μ m), distributing sections progressively across sets of 3-4 slides so that each slide has non-consecutive sections.

Note

Chrome-alum treated slides are pre-coated (subbed) with 1% gelatin, dried and stored at room temperature. This slide treatment enhances section adhesion while retaining good surface tension of the antibody droplets.

Immunostaining

- 3 Air-dry slides at room temperature for at least 10 minutes.
- 4 Draw around sections with hydrophobic barrier pen (PAP pen); wait for 10 min to dry.
- 5 Wash sections in PBS (10 min).
- 6 Incubate sections in blocking solution (PBS containing 10% non-immune horse serum and 0.5% triton X-100) at room temperature in a humidified dark chamber for 2 h.
- 7 Incubate sections in appropriate dilutions of primary antibodies (or combinations of primary antibodies) for 48h in a humidified dark chamber. Antibodies are diluted in PBS containing 2% horse serum, 0.5% Triton and 0.1% sodium azide.
- 8 Wash sections in PBS (30 min)

-
- 9** Incubate sections in appropriate dilutions of secondary antibodies (or combinations of secondary antibodies) for 4h in a humidified dark chamber. Antibodies are diluted in PBS containing 2% horse serum, 0.5% Triton and 0.1% sodium azide.
 - 10** Wash sections in PBS (30 min)
 - 11** Coverslip in Vectorshield or preferred anti-fade mountant.