NOV 17, 2017

Immunohistochemical staining

Xiao Bing Tang¹, Li Huan¹, Jin Zhang¹, Wei Lin Wang¹,
Zheng Wei Yuan², Yu Zuo Bai¹

¹Department of Pediatric Surgery, Shengjing Hospital, China Medical University;

²The Key Laboratory of Health Ministry for Congenital Malformation

Xiaobing Tang

OPEN ACCESS**DOI:**

dx.doi.org/10.17504/protocols.io.kujcwun

Protocol Citation: Xiao Bing Tang, Li Huan, Jin Zhang, Wei Lin Wang, Zheng Wei Yuan, Yu Zuo Bai 2017.

Immunohistochemical staining.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.kujcwun>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Nov 17, 2017

Last Modified: Mar 23, 2018

PROTOCOL integer ID:
8811

- 1 Slides are deparaffinized by immersion in xylene 2 times for 2 hr.

- 2 Slides are rehydrated by immersion in 100% ethanol (10 min.), in 95% ethanol (10 min), in 85% ethanol (10min). Slides are washed with PBS (3 times, 3 min).
- 3 Antigen retrieval is performed by heating the slides in 10 mM citrate buffer (pH 6.0) at 98°C for 10 min in a microwave oven. Slides are washed with PBS (3 times, 3 min).
- 4 The endogenous peroxidase activity was blocked by incubation in 3% H₂O₂ for 20 min. Slides are washed with PBS (3 times, 3 min).
- 5 Antibody incubations were performed in phosphate-buffered saline (PBS) supplemented with 10% goat serum for 20 min at room temperature.
- 6 Slides are incubated 16hr at 4°C in a humidifying box with primary antibody. Negative controls are made with PBS alone.
- 7 Slides are washed with PBS (3 times, 3 min).
- 8 Slides are incubated with biotinylated secondary antibody for 30 min in a humidifying box at room temperature.
- 9 Slides are washed with PBS (3 times, 3 min).

- 10** Slides are incubated with horseradish peroxidase (HRP)-conjugated secondary antibody (Santa Cruz Bio-technology) for 20 min in a humidifying box at room temperature.
- 11** Slides are washed with PBS (3 times, 3 min).
- 12** Signals were visualized by using 3 '3Pdiaminobenzidine (DAB; Sigma, UK) for 1 minute and terminated by incubated in distilled water.
- 13** Slides are washed with distilled water and counterstained with hematoxylin for 10 seconds.
- 14** Slides are washed with running water for 6 min.
- 15** Slides are dehydrated by immersion in 70% ethanol (1 min), in 95% ethanol (5 min), in 100% ethanol (2 times, 5 min) and in xylene (2 times, 20 min).
- 16** Cover slides are mounted with gum.