VERSION 1

SEP 17, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bmgfk3tn

External link:
<https://doi.org/10.1371/journal.pone.0252607>

Protocol Citation: Lora Starrs, Debra Brown, Sarah Popp, Charmaine Simeonovic 2020. Immunohistochemical staining of CD44 core proteins in islet beta cells of formalin-fixed mouse pancreas. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bmgfk3tn>

MANUSCRIPT CITATION:
 Dhouchak S, Popp SK, Brown DJ, Laybutt DR, Biden TJ, Bornstein SR, Parish CR, Simeonovic CJ (2021) Heparan sulfate proteoglycans in beta cells provide a critical link between endoplasmic reticulum stress, oxidative stress and type 2 diabetes. *PLoS ONE* 16(6): e0252607. doi: [10.1371/journal.pone.0252607](https://doi.org/10.1371/journal.pone.0252607)

Immunohistochemical staining of CD44 core proteins in islet beta cells of formalin-fixed mouse pancreas V.1

Lora Starrs¹, Debra Brown¹, Sarah Popp¹, Charmaine Simeonovic¹

¹The Australian National University

Charmaine Simeonovic

ABSTRACT

Paraffin sections (4mm thickness) of formalin-fixed mouse pancreases were treated with heat/citrate buffer for antigen retrieval. CD44 core proteins were detected immunohistochemically using rat anti-mouse CD44 mAb (IM7 mAb; BD Biosciences) or mouse anti-human CD44v3 mAb (CD44v3: R&D Systems), with horseradish peroxidase-conjugated rabbit anti-rat Ig (Dako) or rabbit anti-mouse Ig (Dako). Background staining was checked using the corresponding isotype control Ig instead of the primary antibody. 3-amino-9-ethylcarbazole (AEC) was used as the chromogen. For morphometry, stained sections were imaged using a light microscope with attached camera (Olympus BX41). Image J software with color deconvolution plugin was used for the quantitative analysis of the % of islet area stained.

GUIDELINES

Rat anti-mouse CD44 (IM7) mAb recognises all CD44 isoforms (1-4) and mouse anti-human CD44v3 mAb recognises a CD44 isoform that carries HS side-chains (1-3). In mouse pancreas we have demonstrated that these two mAbs have a similar pattern of intra-islet staining.

References:

1. Bennett KL, Jackson DG, Simon JC, Tanczos E, Peach R, Modrell B, et al. CD44 isoforms containing exon V3 are responsible for the presentation of heparin-binding growth factor. *The Journal of cell biology*. 1995;128(4):687-98.
2. Jackson DG, Bell JI, Dickinson R, Timans J, Shields J, Whittle N. [Proteoglycan forms of the lymphocyte homing receptor CD44 are alternatively spliced variants containing the v3 exon. The Journal of cell biology. 1995;128\(4\):673-85.](#)
3. Martegani MP, Del Prete F, Gasbarri A, Natali PG, Bartolazzi A. [Structural variability of CD44v molecules and reliability of immunodetection of CD44 isoforms using mAbs specific for CD44 variant exon products. Am J Pathol. 1999;154\(1\):291-300.](#)
4. Rops AL, van der Vlag J, Lensen JF, Wijnhoven TJ, van den Heuvel LP, van Kuppevelt TH, et al. [Heparan sulfate proteoglycans in glomerular inflammation. Kidney international. 2004;65\(3\):768-85.](#)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Sep 17, 2020

Last Modified: Sep 17, 2020

PROTOCOL integer ID:
42215

Keywords: CD44,
Immunohistochemistry, Mouse
pancreas

BEFORE START INSTRUCTIONS

Materials:

1. Prepare graded alcohols and xylene for deparaffinizing tissue sections: 2 x xylene (250 ml/slide container), 2 x absolute ethanol (250 ml/slide container), 1 x 90% ethanol (250 ml), 1 x 70% ethanol (250 ml).
2. Prepare acetate buffer components:
(i) 0.1N acetic acid: 290 ml glacial acetic acid in 50 ml deionized water
(ii) 0.1M sodium acetate: 410 mg anhydrous CH_3COONa in 50 ml deionized water.
Prepare 0.1M acetate buffer (pH 5.2) by mixing 10.5 ml 0.1N acetic acid and 39.5 ml 0.1M sodium acetate.
3. Prepare stock solution of 3-amino-9-ethylcarbazole (AEC; chromogen, 8 mg/ml: 40 mg AEC in 5 ml N-N-dimethyl formamide; protect from light and refrigerate at 4°C.
4. Prepare M.O.M. diluent: 200 ml M.O.M. protein concentrate stock solution (M.O.M immunodetection kit) in 2.5 ml phosphate-buffered saline (PBS) for use either as a blocking step to minimize non-specific Ig binding or for diluting antibodies.
5. Mabs and pAbs:
rat anti-mouse CD44 (IM7) mAb, BD Biosciences #553130 or
mouse anti-human CD44v3 mAb, R&D systems BBA11
horseradish peroxidase (HRP)-conjugated rabbit anti-rat Ig, Dako #PO450 (alternative HRP-rabbit anti-rat IgG, Sigma #A5795)
or HRP-rabbit anti-mouse IgG, Invitrogen #31450 (for use with CD44v3 mAb as primary ab)
Purified NA/LE Rat IgG_{2bk}, BD Biosciences #555845 or
Purified Mouse IgG_{2bk}, BD Biosciences #557351
6. Other reagents:
Hydrogen peroxide (30% w/w), Chem-Supply Pty Ltd (Australia) #HA154-500M,
Methanol, Merck # CAS-No. 67-56-1
3Amino-9-ethylcarbazole (AEC), Sigma-Aldrich #A5754
Animal free blocker, Vector Labs #SP-5030
Stock protein concentrate, M.O.M immunodetection kit, Vector Labs # PK-2200
N-N-dimethyl formamide, Sigma #D158550
Glycergel mounting medium, Dako #C0563

1 See Guidelines before starting

- 2 Deparaffinize slides in each xylene for 1 min. rehydrate slides in graded alcohols beginning in absolute ethanol (10 dips)/ container of absolute ethanol), followed by 90% ethanol (10 dips) and 70% ethanol (10 dips). Wash well in running tap water for 5 min.
- 3 Wipe around sections with a tissue, encircle the sections using a diamond pencil and place in a slide container of tap water (250 ml).
- 4 Block endogenous peroxidase activity by incubating sections in 3% hydrogen peroxide in methanol (25 ml 30% H₂O₂ + 225 ml methanol).
- 5 Wash 2 x 2 min in 250 ml phosphate-buffered saline (PBS) followed by wash in running tap water for 5 min.
- 6 Prepare citrate buffer, pH 6 for antigen retrieval. Dissolve 1.05 g Citric acid in 500 ml deionized water and pH using 2 - 10 M NaOH.
- 7 Transfer slides to 250ml citrate buffer and heat in microwave (1600 watt) for 2 min on High power followed by 2 x 6 min on Low power. Allow the slides to cool on the bench for 30 min. Wash slides in 250 ml PBS, 3 x 10 min.
- 8 Wipe around sections using tissue. To block non-specific binding of Ig, apply animal free block (diluted to 20% v/v with deionized water) to tissue sections and incubate for 10 min at room temperature.
- 9 Tip off excess block in Step 7, wipe around sections using tissue and incubate with 40 µg/ml anti-CD44 (IM7) mAb (or 40 µg/ml rat IgG_{2bK} as isotype control; diluted in M.O.M. diluent), 125-150 µl/section at room temperature for 1 hour. Alternatively use 10 µg/ml mouse anti-human CD44v3 mAb (or 10 µg/ml mouse IgG_{2bK} as isotype control Ig)

- 10** Wash off primary antibody with PBS and transfer slides to slide container with 250 ml PBS. Wash 2 x 2min.
- 11** Wipe around sections using tissue and incubate with 52 µg/ml secondary HRP-rabbit anti-mouse Ig (Dako; or alternative 30-60 µg/ml, Sigma) , 150 µl/section, for 30 min at room temperature.
- 12** Wash off secondary antibody with PBS and transfer to slide container with 250 ml PBS. Wash slides 2 x 2min.
- 13** Prepare AEC working solution: 4.75 ml acetate buffer (see Guidelines), 0.25ml AEC stock solution and 25 µl 3% H₂O₂. Filter using a disposable 0.2 µm filter. Use within 2 hours of preparation, refrigerate for short-term storage. Protect from light.
- 14** Wipe around sections using tissue and cover the sections with AEC solution for 30 min at room temperature.
- 15** Wash off AEC solution with deionized water and transfer slides to slide container with 250 ml deionized water. Wash 3x in 10 min.
- 16** Lightly counterstain with Gill's hematoxylin, wash in deionized water (2 x) and briefly dip in ammonium water (100 µl ammonia in 250 ml deionized water), 2 x 2 sec. Wash in deionized water (2x in 250 ml) and coverslip using glycergel mounting medium.
- 17** Image sections using a light microscope with camera attachment. Use Image J software with color deconvolution plugin to determine % of islet area stained.