JUL 12, 2019

Immunohistochemistry- mounted slides

John Tompkins¹, Don Hoover², Jeffrey Ardell¹,
Kalyanam Shivkumar¹

¹University of California, Los Angeles; ²ETSU

Scott John

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.2pagdie

Protocol Citation: John Tompkins, Don Hoover, Jeffrey Ardell, Kalyanam Shivkumar 2019. Immunohistochemistry-mounted slides. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.2pagdie>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: May 07, 2019

Last Modified: Jul 12, 2019

PROTOCOL integer ID:
22978

1 Day 1

Ø Using the PAP Pen, carefully draw a water barrier circle around the tissue sections on the slide
– allow this circle to dry for several seconds or up to approx. one min.

- Ø Rinse slides with PBS (pH 7.3-7.4): 4 x 5 min each
- Ø Rinse slides with 0.5% BSA + 0.4% Triton X-100 in PBS): 1 x 10 min
- Ø Remove slides one at a time and using a clean Kimwipe, carefully wipe around the tissue sections to dry the slide.
- Ø Place the slides into a black, covered slide incubation box/humidity box
- Ø Cover the tissue sections with blocking buffer (10% normal donkey serum in 1.0% BSA + 0.4% Triton X-100 + PBS)
- Ø Allow the sections to remain in blocking buffer for 1.5-2 hrs. at RT
- Ø Pour off the blocking buffer
- Ø Replace with primary antibody solution (antibody of choice diluted in 1.0% BSA + 0.4% Triton X-100 + PBS)
- Ø Incubate tissue with primary antibody overnight in incubation box.

2 Day 2

- ### 3
- Ø Rinse slides with PBS: 4 x 5 min
 - Ø Rinse slides with 0.5% BSA + 0.4% Triton X-100 + PBS: 1 x 10 min
 - Ø Prepare fluorescent secondary antibody (secondary antibody should be diluted in 1.0% BSA + 0.4% Triton X-100 + PBS) (*use Biotin-SP for the antibody that will later add Streptavidin to; if there is a second primary then use the normal secondary for it*).
 - Ø Cover the tissue with the secondary antibody solution and incubate for 2 hrs. at RT in the incubation box. *From this point on, use low light and/or cover tissues.*
 - Ø Rinse slides with PBS: 4 x 5 min
 - Ø Remove excess PBS with a Kimwipe.
 - Ø Prepare fluorescent Streptavidin secondary antibody (should be diluted in 1.0% BSA + 0.4% Triton X-100 + PBS)
 - Ø Cover the tissue with the fluorescent Streptavidin antibody solution and incubate for 2 hrs. at RT in the incubation box.
 - Ø Rinse slides with PBS: 4 x 5 min
 - Ø Remove excess PBS with a Kimwipe.
 - Ø Carefully add a drop of Citifluor (or equivalent) mounting medium to the center of the tissue and apply cover glass.
 - Ø Seal cover glass with clear nail polish. For thicker tissue, add a weight before sealing.