Immunohistochemistry Protocol for Paraffin-Embedded Sections V.2

Kelsey Knight¹

¹BioLegend

Sam Li
BioLegend

DOI:
[dx.doi.org/10.17504/protocol.s.io.tkjekun](https://doi.org/10.17504/protocol.s.io.tkjekun)

External link:
<https://www.biolegend.com/protocols/immunohistochemistry-protocol-for-paraffin-embedded-sections/4256/>

Protocol Citation: Kelsey Knight . Immunohistochemistry Protocol for Paraffin-Embedded Sections.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.tkjekun>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Created: Sep 13, 2018

Last Modified: Aug 29, 2019

PROTOCOL integer ID:
15723

Keywords: IHC

SAFETY WARNINGS

⚠ Formalin is a suspect carcinogen. It can cause eye, skin, and respiratory tract irritation. It should be handled in a hood.

DAB is a suspect carcinogen. Handle with care. Wear gloves, lab coat and eye protection.

BEFORE START INSTRUCTIONS

- For initial experiments, the user must titrate primary and secondary reagents so that staining with the secondary antibody alone yields no background while staining with primary and secondary antibodies yields strong, specific staining.
- Take care to ensure that slides do not dry out by incubating with sufficient volumes and/or in a humidified chamber (such as [926301](#)).

Prepare formalin-fixed, paraffin-embedded tissue sections

- 1 Fix freshly dissected tissue (<3mm thick) with 10% formalin or other fixatives for 24-48 h at room temperature.

🕒 24:00:00

Note

(Caution: Formalin is a suspect carcinogen. It can cause eye, skin, and respiratory tract irritation. It should be handled in a hood.)

- 2 Rinse the tissue with running tap water for 1 h.
01:00:00
- 3 Dehydrate the tissue through 70%, 80%, 95% alcohol, 45 min each, followed by 3 changes of 100% alcohol, 1 hour each.
00:45:00
01:00:00
- 4 Clear the tissue through 2 changes of xylene, 1 hour each.
01:00:00
- 5 Immerse the tissue in 3 changes of paraffin, 1 hour each.
01:00:00
- 6 Embed the tissue in a paraffin block. The paraffin tissue block can be stored at room temperature for years.
- 7 Section the paraffin-embedded tissue block at 5-8 μ m thickness on a microtome and float in a 40°C water bath containing distilled water.
- 8 Transfer the sections onto glass slides suitable for immunohistochemistry (e.g. Superfrost Plus). Allow the slides to dry overnight and store slides at room temperature until ready for use.

- 9 Deparaffinize slides in 2 changes of xylene, 5 min each.
00:05:00
- 10 Transfer slides to 100% alcohol, for 2 changes, 3 min each, and then transfer once through 95%, 70% and 50% alcohols respectively for 3 min each. 00:03:00
- 11 Block endogenous peroxidase activity by incubating sections in 3% H₂O₂ solution in methanol at room temperature for 10 min to block endogenous peroxidase activity.
00:10:00
- 12 Rinse in 300ml of PBS for 2 changes, 5 min each.
00:05:00
- 13 *Optional:* Perform antigen retrieval to unmask the antigenic epitope. The most commonly used antigen retrieval is a citrate buffer method. Arrange the slides in a staining container. Pour 300ml of 10mM citrate buffer, pH 6.0 into the staining container and incubate it at 95-100°C for 10 min (optimal incubation time should be determined by user). Remove the staining container to room temperature and allow the slides to cool for 20 min.
- 14 Rinse slides in 300ml PBS for 2 changes, 5 min each.
- 15 *Optional:* Add 100µl blocking buffer (e.g. 10% fetal bovine serum in PBS) onto the sections of the slides and incubate in a humidified chamber at room temperature for 1 hour. 01:00:00
- 16 Drain off the blocking buffer from the slides.
- 17 Apply 100µl appropriately diluted primary antibody (in antibody dilution buffer, e.g. 0.5% bovine serum albumin in PBS) to the sections on the slides and incubate in a humidified chamber at room temperature for 1 hour. 01:00:00

- 18** Wash the slides in 300ml PBS for 2 changes 5 min each.
00:05:00
- 19** Apply 100µl appropriately diluted biotinylated secondary antibody (using the antibody dilution buffer) to the sections on the slides and incubate in a humidified chamber at room temperature for 30 min. 00:30:00
- 20** Wash slides in 300ml PBS for 2 changes, 5 min each. 00:05:00
- 21** Apply 100µl appropriately diluted Sav-HRP conjugates (using the antibody dilution buffer) to the sections on the slides and incubate in a humidified chamber at room temperature for 30 min (keep protected from light).
00:30:00
- 22** Wash slides in 300ml PBS for 2 changes, 5 min each.
00:05:00
- 23** Apply 100 µl DAB substrate solution (freshly made just before use: 0.05% DAB - 0.015% H₂O₂ in PBS) to the sections on the slides to reveal the color of antibody staining. Allow the color development for < 5 min until the desired color intensity is reached.
- Note**
- (Caution: DAB is a suspect carcinogen. Handle with care. Wear gloves, lab coat and eye protection.)
- 24** Wash slides in 300ml PBS for 2 changes, 5 min each.
00:02:00
- 25** *Optional:* Counterstain slides by immersing sides in Hematoxylin (e.g. Gill's Hematoxylin) for 1-2 min.
00:02:00
- 26** Rinse the slides in running tap water for > 15 min.

00:15:00

- 27** Dehydrate the tissue slides through 4 changes of alcohol (95%, 95%, 100% and 100%), 5 min each.

00:05:00

- 28** Clear the tissue slides in 3 changes of xylene and coverslip using mounting solution (e.g. Permount). The mounted slides can be stored at room temperature permanently.

- 29** Observe the color of the antibody staining in the tissue sections under microscopy.