Immunochemistry Protocol for Sternberger Monoclonal Antibodies V.2

Kelsey Miller¹

¹BioLegend

Sam Li
BioLegend

GUIDELINES

Use with Ultra Streptavidin Detection Kit ([SIG-32250](#)) or ([SIG-32248](#))

Positive control: Normal human cerebellum (except SMI-71, which should be rat brain)

DOI:
[dx.doi.org/10.17504/protocols.io.tkkekuw](https://doi.org/10.17504/protocols.io.tkkekuw)

External link:
<https://www.biolegend.com/protocols/immunohistochemistry-protocol-for-sternberger-monoclonal-antibodies/4253/>

Protocol Citation: Kelsey Miller . Immunohistochemistry Protocol for Sternberger Monoclonal Antibodies.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.tkkekuw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Created: Sep 13, 2018

Last Modified: Aug 29, 2019

PROTOCOL integer ID:
15724

Keywords: IHC

Clear Slides

1 Clear Slides: Removes paraffin and hydrates the tissue.

Xylene	5 minutes in each of (3) different 250mL containers
100% alcohol	5 minutes in each of (3) different 250mL containers

95% alcohol	3 minutes in (1) 250mL container
70% alcohol	3 minutes in (1) 250mL container
water	1 minutes in each of (3) different 250mL containers
H2O2 (3%)	15 minutes in (1) 250mL container

Rinse slides

- 2 Rinse slides with lab grade water.

Note: Lab grade filtered water such as injection grade, cell culture grade, Reverse Osmosis De-ionization (RODI)

Antigen Retrieval

- 3 Heat slides in 1X Retrieve ALL3 solution for 1 minute 40 seconds on high power in microwave

00:01:40

- 4 Reduce to low power and simmer 10 minutes in microwave.

00:10:00

- 5 Remove from microwave and allow slides to cool on the bench top for 10 minutes.

00:10:00

- 6 Rinse slides with lab grade water.

- 7 Apply serum block for at least 5 minutes. Do NOT wash after this step.

00:05:00

- 8 Blot off serum block.

- 9 Apply primary antibody (see recommended dilution from datasheet).
- 10 Incubate primary antibody 60 minutes at room temperature.
01:00:00
- 11 Rinse slides with 1X PBS.
- 12 Apply USA Linking reagent - 20 minutes incubation.
(If using a biotinylated product, omit this step)
00:20:00
- 13 Rinse slides with 1X PBS.
- 14 Apply Labeling Reagent - 20 minutes incubation. 00:20:00
- 15 Rinse with 1X PBS.
- 16 Apply chromogen - 5 minutes incubation. Dilute according to manufacturer's instructions.
00:05:00
1. AEC Chromogen: 20µL AEC chromogen + 1mL AEC substrate buffer
 2. DAB Chromogen: 40µL DAB chromogen + 1mL DAB substrate buffer

Counterstain

17 Submerge slides in Mayer's Hematoxylin for 30 seconds.

00:00:30

18 Rinse under running lab grade water for 1 minute or until water is clear.

00:01:00

19 Submerge slides in Bluing Reagent for 1 minute.

00:01:00

20 Rinse under running lab grade water for 1 minute.

00:01:00

Clear slides

21 Clear slides: Dehydrate the tissue.

1. 95% alcohol 3 minutes in (1) 250mL container
2. 100% alcohol 5 minutes in each of (3) different 250mL container
3. Xylene 5 minutes in each of (3) different 250mL container

Coverslip

22 Cover slip slide using Permanent Aqueous Mounting Medium ([SIG-31010](#)).

Note: Do not use xylene based mount with AEC Chromogen as it will dissolve the chromogen