



Oct 24, 2023

Version 2

Improving genome-wide mapping of nucleosomes in *Trypanosome cruzi*. V.2

 PLOS One

 Peer-reviewed method

DOI

dx.doi.org/10.17504/protocols.io.6qpvr4w83gmk/v2

Paula Beati¹, Milena Massimino Stepñicka¹, Salomé C. Vilchez Larrea^{1,2}, Pablo Smircich^{3,4}, Guillermo Daniel Alonso^{1,2}, Josefina Ocampo¹

¹Instituto de Investigaciones en Ingeniería Genética y Biología Molecular, Consejo Nacional de Investigaciones Científicas y Técnicas, Vuelta de Obligado 2490, C1428ADN Buenos Aires, Argentina;

²Departamento de Fisiología, Biología Molecular y Celular, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Buenos Aires, Argentina;

³Laboratorio de Bioinformática, Departamento de Genómica, Instituto de Investigaciones Biológicas Clemente Estable (IIBCE), Montevideo, Uruguay;

⁴Sección Genómica Funcional, Facultad de Ciencias, Universidad de la República (Udelar), Montevideo, Uruguay.

Milena Massimino Stepñicka: Former member;

PLOS ONE Lab Protocols

Tech. support email: plosone@plos.org



Josefina Ocampo

INGEBI CONICET

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account





DOI: <https://dx.doi.org/10.17504/protocols.io.6qpvr4w83gmk/v2>

External link: <https://doi.org/10.1371/journal.pone.0293809>

Protocol Citation: Paula Beati, Milena Massimino Stepñicka, Salomé C. Vilchez Larrea, Pablo Smircich, Guillermo Daniel Alonso, Josefina Ocampo 2023. Improving genome-wide mapping of nucleosomes in Trypanosome cruzi.. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.6qpvr4w83gmk/v2> Version created by **Josefina Ocampo**

Manuscript citation:

Beati P, Massimino Stepñicka M, Vilchez Larrea SC, Smircich P, Alonso GD, et al. (2023) Improving genome-wide mapping of nucleosomes in *Trypanosome cruzi*. PLOS ONE 18(11): e0293809. <https://doi.org/10.1371/journal.pone.0293809>

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: October 18, 2023

Last Modified: October 24, 2023

Protocol Integer ID: 89570

Keywords: Chromatin; Nucleosomes; Micrococcal nuclease; genome-wide mapping; Trypanosoma cruzi, nucleosomes genome, whole genome refines the average nucleosome representation, genes in trypanosome, wide mapping of nucleosome, uniqueness of the chromatin organization, chromatin organization, determining nucleosome position, average nucleosome representation, nucleosome position, nucleosome, chromatin, histone protein, intrypanosoma cruzi dna, chromatin by octamer, digestion of chromatin, trypanosome, trypanosome cruzi, complex genome, genome, improving genome, whole genome, whole genome refine, transcription of protein, coding gene, gene expression, micrococcal nuclease, transcription, gene, deep sequencing, hybrid reference strain, protein, dna, dna dependent process, spurious alignment, method to the hybrid reference strain

Funders Acknowledgements:

Agencia Nacional de Promoción Científica y Tecnológica

Grant ID: PICT 2015-0898

Agencia Nacional de Promoción Científica y Tecnológica

Grant ID: PICT 2018-00621

Agencia Nacional de Promoción Científica y Tecnológica

Grant ID: PICT 2020-00473

Abstract

In *Trypanosoma cruzi* DNA is packaged into chromatin by octamers of histone proteins that form nucleosomes. Transcription of protein coding genes in trypanosomes is constitutive producing polycistronic units and gene expression is primarily regulated post-transcriptionally. However, chromatin organization influences DNA dependent processes. Hence, determining nucleosome position is of uppermost importance to understand the peculiarities found in trypanosomes. To map nucleosomes genome-wide in several organisms, digestion of chromatin with micrococcal nuclease followed by deep sequencing has been applied. Nonetheless, the special requirements for cell manipulation and the uniqueness of the chromatin organization in trypanosomes entails a customized analytical approach. In this work, we adjusted this broadly used method to the hybrid reference strain, CL Brener. Particularly, we implemented an exhaustive and thorough computational workflow to overcome the difficulties imposed by this complex genome. We tested the performance of two aligners, Bowtie2 and HISAT2, and discuss their advantages and caveats. Specifically, we highlight the relevance of using the whole genome as a reference instead of the commonly used Esmeraldo-like haplotype to avoid spurious alignments. Moreover, we show that using the whole genome refines the average nucleosome representation, but also the quality of mapping for every region represented. Furthermore, we provide a source code for the construction of 2D plots and heatmaps which are easy to adapt to any *T. cruzi* strain.

Guidelines

Do not vortex at any step of the protocol.

Add protease inhibitors fresh.

The range of MNase to be tested is only a suggestion. We recommend performing a titration in every replicate experiment.

Materials

Special material for MNase digestion

Micrococcal Nuclease (MNase), Worthington Biochemical Corporation, Cat No: R1A12424

Special material for library preparation

PrePCR kit (New England Biolabs, Ipswich, MA, US)

Qiagen PCR column (Qiagen, Hilden, Germany)

NEBNext Ultra DNA Library Prep Kit for Illumina and NEBNext Multiplex Oligos for Illumina (New England Biolabs, Ipswich, MA, US).

Agencourt AMPure XP beads (Beckman-Coulter, Brea, CA, US)



Troubleshooting

Before start

Prepare every solution in advance before you start with the MNase digestion.

Incomplete lysis solution

(1 mM L-glutamine, 250 mM sucrose, 2.5 mM CaCl_2 , 1 mM phenylmethylsulfonyl fluoride (PMSF)).

Complete lysis Solution

(10 ml incomplete Lysis Solution, 0.01% Triton X 100, 1 mM PMSF).

MNase digestion buffer

(10 mM HEPES, 35 mM NaCl, 500 μM MgCl_2 , 500 μM CaCl_2 , 5 mM 2-mercaptoethanol and protease inhibitors)

MNase stock solution

Disolve Micrococcal Nuclease (MNase), Worthington Biochemical Corporation, Cat No: R1A12424 25,271 U/mg P, 82% protein, 45 Ku in 4.5 ml MNase storage buffer, aliquot out in 9× 500ml. Stored at -80°C .


MNase storage buffer

(5 mM Na-phosphate, pH7.0, 0.025mM CaCl_2)



MNase Stop Solution

(180 mM EDTA, 7.2 % SDS prepared in H_2O)







Epimastigote cultivation and harvest

- 1 CL Brener epimastigotes were grown at  28 °C in liver infusion tryptose (LIT) medium (5 g.l-1 liver infusion, 5 g.l-1 bacto-tryptose, 68 mM NaCl, 5.3 mM KCl, 22 mM Na₂HPO₄, 0.2% (w/v) glucose, and 0.002 % (w/v) hemin) supplemented with 10 % (v/v) fetal calf serum (FCS), 100 U/ml penicillin and 100 mg. l⁻¹ streptomycin. Cell density was maintained between 1×10⁶ and 1×10⁸ cells.ml⁻¹ by sub-culturing parasites every 6-7 días.


MNase digestion of *T. cruzi* epimastigote chromatin

- 2 Pellet 10e⁸ late exponentially growing epimastigotes per reaction (normally 7) by centrifugation at 1700 xg for 5 minutes at  Room temperature . Wash the cells with 10 ml Incomplete Lysis Solution (1 mM L-glutamine, 250 mM sucrose, 2.5 mM CaCl₂, 1 mM phenylmethylsulfonyl fluoride (PMSF)).
- 3 Centrifuge at 1700 xg for 5 minutes and discard supernatant.
- 4 Permeabilize cells with 800 µl Complete Lysis Solution (10 ml Incomplete Lysis Solution, 0.01% Triton X 100, 1 mM PMSF).
- 5 Divide the material into 7 aliquots of 100 µl each into 1.5 ml microcentrifuge tubes.
- 6 Centrifuge at 1700 xg for 5 minutes and discard supernatant.
- 7 Wash the pellets with 100 µl Incomplete lysis solution.
- 8 Centrifuge at 1700 xg por 5 minutos.
- 9 Resuspend the pellets in MNase Digestion Buffer (10 mM HEPES, 35 mM NaCl, 500 µM MgCl₂, 500 µM CaCl₂, 5 mM 2-mercaptoethanol and protease inhibitors). Pre-warm the buffer in a  25 °C water bath for 2 min before resuspension



- 10 Add increasing amounts of MNase (10 U/μl) to each tube, for example: 0, 1, 3.5, 7, 10, 20, 40 μl, and mix thoroughly without vortexing. Incubate the digests at  25 °C for 5 min. Stop the reaction by adding 50 μl MNase Stop Solution (180 mM EDTA, 7.2 % SDS prepared in H₂O) to each digestion, mixing, and leaving at room temperature for 5 min. 
- 11 Add 50 μl of 10 % SDS and mix (final SDS ~1.8% in ~500 μl final volume).
- 12 Add 250 μl of 3M KOAc (~1 M final in ~750 μl final volume) and mix.
- 13 Extract the DNA by adding 750 μl chloroform, mixing and spinning in a microcentrifuge (5min at top speed). Repeat extraction.
- 14 Transfer the supernatants to new 1.5-ml tubes and add 0.7 vol. isopropanol to precipitate the DNA. Mix and leave at  -20 °C  Overnight 
- 15 Pellet the DNA by spinning in a microcentrifuge (30 min at top speed).
- 16 Discard the supernatant. Wash the pellet once with 0.6 ml 70 % EtOH and spin in a microcentrifuge (5 min at top speed).
- 17 Discard the supernatant and dissolve each pellet in 50 μl of 10 mM Tris HCl pH 8, 2 mM EDTA with 100 μg/ml of RNaseA. Incubate the samples at  37 °C for 2 hours.

Paired-end sequencing

- 18 Paired-end libraries were prepared as described before (1,2). 
Briefly, once the DNA samples with the appropriate level of digestion were selected, any nicks were repaired using the PrePCR kit (New England Biolabs, Ipswich, MA, US), purified using a Qiagen PCR column (Qiagen, Hilden, Germany) and eluted with 50 μl TE buffer (10 mM Tris-Cl pH8, 0.1 mM EDTA). Modification of the 5' and 3' ends of the nucleosomal DNA and adapter ligation were performed with the NEBNext Ultra DNA Library Prep Kit for Illumina and NEBNext Multiplex Oligos for Illumina (New England Biolabs, Ipswich, MA, US). Agencourt AMPure XP beads (Beckman-Coulter, Brea, CA, US) were used to purify the DNA after adaptor ligation and PCR products before sequencing as described (1,2). The beads were used in a 1:1 volume ratio,

which removes adaptors and any primer dimers but retains nucleosomal DNA ligated to adaptors and anything larger. For sequencing reactions, either Illumina Next-seq 500 or Hi-seq 2000 were used to obtain pairs of 50 nucleotide (nt) reads.

Data files and statistical analysis

- 19 Paired-end reads are normally provided in the fastq file format containing the raw reads. To do quality control checks on raw data, fastQC analysis was performed. The fastQC reports of the current data could be found at https://github.com/paulati/nucleosome/tree/master/output/qc_output. During this step over-represented sequences (ORS) were detected and trimmed out to avoid the generation of artefacts during the analysis. ORS in general are 50 nts long, when working with 50 nts reads, and correspond to adaptors or primers used for library construction as detailed in the fastQC report. In the case of *T. cruzi*, due to the repetitive nature of its genome, some ORS occasionally contain runs of repetitive sequences. ORS were removed when present producing trimmed fastq files using the Cutadapt tool (3), containing trimmed raw reads that from now on we will refer to as "*trimmed reads*". Given that this step is regularly omitted when analyzing MNase-seq data from model organisms, to evaluate if it constitutes an improvement in the analysis of *T. cruzi*, we pursued the following steps in parallel using raw reads and trimmed reads.



Sequence alignment and reference genome

- 20 Bowtie2 (4) and HISAT2 (5) were used to align either the raw or the trimmed 50 nt paired-end reads, to the corresponding reference genome. To compare the performance of these tools, the same parameter values were set for both tools. In particular, both tools have a parameter called score-min, defined as a function of the read length, that is the minimum score needed for an alignment to be considered "valid".

For Bowtie2, the end-to-end mode was used with default settings, commonly used for model organisms, where the score-min is specified by ``L, -0.6, -0.6``. This establishes a linear function $f(x) = -0.6 + -0.6 * x$, where x is the read length. In this case, x is 50 nts, which implies it accepts scores bigger than -30.6. In the case of HISAT2, the default settings are more restrictive than those for Bowtie2 with ``L, 0, -0.2``. To make a fair comparison between both tools, the exact same score-min used for Bowtie2 was set for HISAT2.

Esmeraldo-like is considered the reference genome. In this work, this was used as a reference and two additional reference genomes were generated for comparison. One, combining Esmeraldo-like and non-Esmeraldo-like haplotypes, designated *Es_U_nonEs*. The second one, combining Esmeraldo-like, non-Esmeraldo-like and "extra" regions that are still not assigned to any haplotype and are annotated separately. This wider genome, is designated *Cl Brener_all*. Versions TriTryp46 of the respective genomes were used as a reference for informatic analysis in every case.



Binary file generation

- 21 Bigwig files containing information of nucleosome occupancy and nucleosome position were generated. Occupancy (coverage) maps were built by counting the number of times that a base pair was occupied by a nucleosome, while position maps counted the number of times each particular nucleosome sequence was present in the data set. The midpoint of the nucleosome, called the dyad, was represented as previously described (6–8). To observe more clearly the nucleosomes in the occupancy maps and to be more accurate in the estimation of the position maps, the analysis was restricted to those fragments that belong to the nucleosome size range (120–180 bp) and is normalized by summing all the nucleosome sequences covering a nucleotide and dividing that number by the average number of nucleosome sequences per base pair across the genome.



Prediction of the trans-splicing acceptor site

- 22 The 5' untranslated regions (5'UTR) were predicted with UTRme (9). Given the lack of transcriptomic data for CL Brener epimastigotes, the predictions were based on available RNA-seq data for the Y strain (10) using TriTryp46 Esmeraldo-like genome as a reference. As an approximation of the trans-splicing acceptor site (TAS), the 5' end of the 5'UTR region predicted with the best score were used.



Average plots and 2D occupancy plots

- 23 Average representation of the data and 2D Occupancy plots were performed as previously described (11,12), with some modifications. Average nucleosome determinations were performed from the alignments generated in all the alternative alignments and represented relative to the predicted TAS in a 1kb window. Briefly, even when the total number of reads was previously aligned to the *Es_U_nonEsor* the *CL Brener_all* genomes, for this analysis only the average nucleosome occupancy coming from the reads that aligned to the Esmeraldo-like haplotype was represented relative to the predicted TAS. We define nucleosome density as the normalized percentage of occupancy. For 2D Occupancy plots, the nucleosome density was represented relative to the TAS in the x axis, while the size of the analyzed DNA fragments was represented in the y axis. For heatmaps showing every region around TAS, nucleosome density was represented in the same way in the x axis; while in the y axis, for each gene region, we computed the normalized total occupancy for the region as the normalized sum of occupancies for each position in the region, and then ordered the regions according to these descending values.



Protocol references

- 1.Ocampo J, Chereji R V, Eriksson PR, Clark DJ. The ISW1 and CHD1 ATP-dependent chromatin remodelers compete to set nucleosome spacing in vivo. *Nucleic Acids Res.* 2016;44(10):4625–35.
- 2.Ocampo J, Chereji R V, Eriksson PR, Clark DJ. Contrasting roles of the RSC and ISW1/CHD1 chromatin remodelers in RNA polymerase II elongation and termination. *Genome Res [Internet]*. 2019 Mar;29(3):407–17. Available from: <http://genome.cshlp.org/lookup/doi/10.1101/gr.242032.118>
- 3.Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal [Internet]*. 2011 May 2;17(1):10. Available from: <http://journal.embnet.org/index.php/embnetjournal/article/view/200>
- 4.Langmead and Steven L Salzberg. Bowtie2. *Nat Methods [Internet]*. 2013;9(4):357–9. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3322381/pdf/nihms-366740.pdf>
- 5.Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory requirements Daehwan HHS Public Access. *Nat Methods.* 2015;12(4):357–60.
- 6.Clark DJ. Nucleosome positioning, nucleosome spacing and the nucleosome code. *J Biomol Struct Dyn [Internet]*. 2010;27(6):781–93. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2935628&tool=pmcentrez&rendertype=abstract>
- 7.Cole HA, Howard BH, Clark DJ. Genome-wide mapping of nucleosomes in yeast using paired-end sequencing [Internet]. 1st ed. Vol. 513, *Methods in Enzymology*. Elsevier Inc.; 2012. 145–168 p. Available from: <http://dx.doi.org/10.1016/B978-0-12-391938-0.00006-9>
- 8.Chereji R V, Clark DJ. Major Determinants of Nucleosome Positioning. *Biophys J [Internet]*. 2018;114(10):1–11. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S0006349518303813>
- 9.Radío S, Fort RS, Garat B, Sotelo-Silveira J, Smircich P. UTRme: A Scoring-Based Tool to Annotate Untranslated Regions in Trypanosomatid Genomes. *Front Genet.* 2018;9(December):1–9.
- 10.Li Y, Shah-Simpson S, Okrah K, Belew AT, Choi J, Caradonna KL, et al. Transcriptome Remodeling in *Trypanosoma cruzi* and Human Cells during Intracellular Infection. *PLoS Pathog [Internet]*. 2016 Apr;12(4):e1005511. Available from: <http://dx.doi.org/10.1371/journal.ppat.1005511>
- 11.Chereji R V, Ocampo J, Clark DJ. MNase-Sensitive Complexes in Yeast: Nucleosomes and Non-histone Barriers. *Mol Cell.* 2017;65(3):565–577.e3.
- 12.Beati P, Chereji R V. Creating 2D Occupancy Plots Using plot2DO. In: *Methods in molecular biology* (Clifton, NJ) [Internet]. 2020. p. 93–108. Available from: <http://link.springer.com/10.1007/978-1-4939-0512-6>