SEP 05, 2018

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.s9qeh5w

Document Citation: Maira Rubi Segura Campos 2018. In vitro α -glucosidase inhibitory assay. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.s9qeh5w>

MANUSCRIPT CITATION:
 Dineshkumar B, Mitra A, Manjunatha M. Studies on the anti-diabetic and hypolipidemic potentials of mangiferin (xanthone glucoside) in streptozotocin-induced type 1 and type 2 diabetic model rats. *Int J Adv Pharm Sci*. 2010; 1: 75-85.

License: This is an open access document distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Created: Sep 05, 2018

Last Modified: Sep 05, 2018

DOCUMENT integer ID:
 15376

In vitro α -glucosidase inhibitory assay

Maira Rubi Segura Campos¹

¹Universidd Aut3noma de Yucat3n

Maira Rubi Segura Campos

ABSTRACT

The *in vitro* α -glucosidase inhibitory assay was performed according to Dineshkumar et al. [8]. The α -glucosidase enzyme (2 U/mL) was premixed with 20 μ L of plant aqueous extract at a concentration of 1,000 μ g/mL and incubated for 5 min at 37 $^{\circ}$ C. Then 1 mM p-nitrophenyl gluco-pyranoside (pNPG) (20 μ L) in 50 mM of phosphate buffer (pH 6.8) was added to initiate the reaction. The mixture was incubated at 37 $^{\circ}$ C for 20 min. The reaction was terminated by the addition of 50 μ L of 1 mM sodium carbonate. The α -glucosidase activity was determined at 405 nm. The α -glucosidase inhibitory activity was calculated using the formula $(Ac^{+}) - (Ac^{-}) - (As - Ab)/(Ac^{+}) - (Ac^{-}) \times 100$, where Ac^{+} , Ac^{-} , As, Ab are defined as the absorbance (405 nm) of 100% enzyme activity (only solvent with enzyme), 0% enzyme activity (only solvent without enzyme), test sample (with enzyme), and a blank (a test sample without enzyme), respectively. Reagents to evaluate anti-diabetic activity were of analytical grade and purchased from Sigma-Aldrich. For anti-diabetic activity, acarbose (anti-diabetic drug) was used as control.